



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/29, 15/63, A01H 5/10</b>		A1	(11) International Publication Number: <b>WO 98/35044</b>
			(43) International Publication Date: 13 August 1998 (13.08.98)
(21) International Application Number: PCT/CA98/00096 (22) International Filing Date: 9 February 1998 (09.02.98) (30) Priority Data: 60/038,815 10 February 1997 (10.02.97) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 60/038,815 (CIP) Filed on 10 February 1997 (10.02.97) (71) Applicant (for all designated States except US): NATIONAL RESEARCH COUNCIL OF CANADA [CA/CA]; 1200 Montreal Road, Ottawa, Ontario K1A 0R6 (CA). (72) Inventors; and (75) Inventors/Applicants (for US only): TAYLOR, David, C. [CA/CA]; 622 Wollaston Bay, Saskatoon, Saskatchewan S7J 4C3 (CA). ZOU, Jitao [CA/CA]; 1619 Acadia Drive, Saskatoon, Saskatchewan S7H 5K7 (CA). (74) Agents: GALE, Edwin, J. et al.; Kirby, Eades, Gale, Baker, Box 3432, Station D, Ottawa, Ontario K1P 6N9 (CA).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

(54) Title: PLANT PYRUVATE DEHYDROGENASE KINASE GENE

## (57) Abstract

The present invention relates to the isolation, purification, characterization and use of a mitochondrial pyruvate dehydrogenase kinase (PDHK) gene [SEQ ID NO:1] (pYA5; ATCC No 209562) from the *Brassicaceae* (specifically *Arabidopsis thaliana*). The invention includes isolated and purified DNA of the stated sequence and relates to methods of regulating fatty acid synthesis, seed oil content, seed size/weight, flowering time, vegetative growth, respiration rate and generation time using the gene and to tissues and plants transformed with the gene. The invention also relates to transgenic plants, plant tissues and plant seeds having a genome containing an introduced DNA sequence of SEQ ID NO:1; or a part of SEQ ID NO:1 characterized in that said sequence has been introduced in an antisense or sense orientation, and a method of producing such plants and plant seeds. The invention also relates to substantially homologous DNA sequences from plants encoding proteins with deduced amino acid sequences of 25 % or greater identity, and 50 % or greater similarity, isolated and/or characterized by known methods using the sequence information of SEQ ID NO:1, and to parts of reduced length that are still able to function as inhibitors of gene expression by use in an antisense, co-suppression or other gene silencing technologies.

```

1: t ooa tct gcy cao tto ttt cgt cca gto gat gat aat aac ggt gg
46: a gaa cga cgg agg cgg cgc acg tta ggg ttt cta atc att tct ct
91: c tct tag acg ctt atg gca gtg aag aaa ggc tgc gaa atg tto co
    M A V X K A C E M F P
136: g aag agt ttg ato gaa gat gtt cao aaa tgg ggt tgc atg aag ca
    K S L I E D V H K W G C M K Q
181: a aoo ggt gtt ago ctt aga tnc atg atg gag ttt ggt tcc aaa cc
    T G V S L R Y M M E F G S X P
226: t act gag agg aat ctt ttg att tct got cag ttt ttg cat aag ga
    T E R N L L I S A Q F L H K E
271: g ctt ccg att cgc gto gcc agg aga gog ato gaa cto cag acg ct
    L P I R V A R R A I E L Q T L
316: t ctt tat ggt cto tct gat aaa cct gcc gtt ttg aag gtg cgg ga
    P Y G L S D K P A V L K V R D
361: t tgg tat ttg gaa tct tto egg gac atg aga gaa ttt cct gag at
    W Y L E S F R D M R A F P E I
406: t aag gat tgg ggt gac gag aaa gat tto act cag atg att aag go
    K D S G D E K D P T Q M I K A
451: t gto aaa gta agy cat aac aat gtg gtt ccc atg atg gct ttg gy
    V K V R N M N V V P M M A L G
496: t gtt aat cag cto aag aaa gga atg aat tot gga aat ctt gat ga
    V M Q L K K G M N S G M L D E
541: g att cat cag ttt ctt gat cgt tto tnc ttg tog oga ato ggy at
    I H Q F L D R F Y L S R I G I
586: c cgg atg ctt att ggg cag cao gtt gag ttg cat aat cca aat co
    R M L I G Q H V S L H N P N P
631: a ccg ctt cat aca gtg ggt tat ata oac aca aag atg tot cct at
    P L H T V G Y I N T K M S P M
676: g gag gta gca agy aat gca agt gaa gat gct cgg tca att tgt tt
    E V A R N A S E D A R S I C F
721: c cga gag taa ggt tot gaa ccg gaa ata aac ata tat ggc gat co
    R E Y G S A P E I N I Y G D P
766: c agt tto aoo ttt ccg tat gtt cca aag cat ttg gat ctt atg at
    S F T F P Y V P T H L D L M M
811: g tat gag cta gto aag aac tot cta cgt gat gto caa gag cga tt
    Y E L V K N S L R A V Q E R F
856: t gtt gac tct gat aga gtt gaa ooa ooa ato cgc att ata gtt go
    V D S D R V A P P I R I I V A

```

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## PLANT PYRUVATE DEHYDROGENASE KINASE GENE

### TECHNICAL FIELD

5

This invention relates to plant genes useful for the genetic manipulation of plant characteristics. More specifically, the invention relates to the identification, isolation and introduction of genes useful, for example, for altering the seed oil content, seed size, flowering and/or generation time, 10 or vegetative growth of commercial or crop plants.

### BACKGROUND ART

Through a coordination of the light and dark reactions of 15 photosynthesis, plants assimilate CO<sub>2</sub> in the formation of sugars. Via the catabolic and anabolic reactions of metabolism, these sugars are the basis of plant growth, and ultimately plant productivity. In the process of plant growth, respiration, which involves the consumption of O<sub>2</sub> and catabolism of sugar or other substrates to produce CO<sub>2</sub>, plays a central role in providing a 20 source of energy, reducing equivalents and an array of intermediates (carbon skeletons) as the building blocks for many essential biosynthetic processes. It is known that any two plants with equal photosynthetic rates often differ in both total biomass production and harvestable product.

Therefore, the relationship between rate of respiration and crop productivity 25 has been one of the most intensively studied topics in plant physiology. In a biochemical sense, respiration can be taken to be composed of glycolysis, the oxidative pentose phosphate pathway, the Krebs' (tricarboxylic acid, TCA) cycle and the mitochondrial electron transport system. The intermediate products of respiration are necessary for growth in meristematic 30 tissues, maintenance of existing phytomass, uptake of nutrients, and intra-

and inter-cellular transport of organic and inorganic materials. In soybean there is evidence that an increase in respiration rate by the pod can lead to an increase in seed growth (Sinclair et al., 1987), while decreased respiration can result in decreased reproductive growth (Gale, 1974). Respiration is therefore important to both anabolic and catabolic phases of metabolism.

Although the pathways of carbon metabolism in plant cells are quite well known, control of the flux of carbon through these pathways *in vivo* is poorly understood at present. The mitochondrial pyruvate dehydrogenase complex (mtPDC), which catalyzes the oxidative decarboxylation of pyruvate to give acetyl CoA, is the primary entry point of carbohydrates into the Krebs cycle. The mtPDC complex links glycolytic carbon metabolism with the Krebs cycle, and, because of the irreversible nature of this reaction, the pyruvate dehydrogenase complex (PDC) is a particularly important site for regulation.

15 Mitochondrial PDC has been studied intensively in mammalian systems, and available knowledge about the molecular structure of plant mtPDC is largely based on studies of the mammalian mtPDC. The mtPDC contains the enzymes E1 (EC 1.2.4.1), E2 (EC 2.3.1.12) and E3 (EC 1.8.1.4) and their associated prosthetic groups, thiamine PPi, lipoic acid, and FAD, respectively. The E1 and E3 components are arranged around a core of E2. The E2 and E3 components are single polypeptide chains. In contrast, the E1 enzyme consists of two subunits, E1 $\alpha$  and E1 $\beta$ . Their precise roles remain unclear. Another subunit, the E3-binding protein, is thought to play a role in attaching E3 to the E2 core. The E1 kinase and phosphatase are 20 associated regulatory subunits (Grof et al., 1995).

Plants are unique in having PDH complexes in two isoforms, one located in the mitochondrial matrix as in other eukaryotic cells, and another located in the chloroplast or plastid stroma (Randall et al., 1989). Although both plastidial and mitochondrial PDH complex isoforms are very sensitive to 30 product feedback regulation, only the mitochondrial PDH complex is

regulated through inactivation/reactivation by reversible phosphorylation/dephosphorylation (Miernyk and Randall, 1987; Gemel and Randall, 1992; Grof et al., 1995). More specifically, the activity of mitochondrial PDC (mtPDC) is regulated through product feedback inhibition (NADH and acetyl-5 CoA) and the phosphorylation state of mtPDC is determined by the combined action of reversible phosphorylation of the E1 $\alpha$  subunit by PDC kinase (PDCK) and its dephosphorylation by PDC phosphatase. PDCK phosphorylates and inactivates PDC, while PDC phosphatase dephosphorylates and reactivates the complex. Maximum PDC activity also 10 appears to vary developmentally, with the highest catalytic activity observed during seed germination and early seedling development (e.g. in post-germinative cotyledons, Hill et al., 1992; Grof et al., 1995).

Acetyl-CoA, the product of PDC, is also the primary substrate for fatty acid synthesis. While it is known that plant fatty acid biosynthesis occurs in 15 plastids, the origin of the acetyl-CoA used for the synthesis of fatty acids in plastids has been the subject of much speculation. It remains a major question which has not been resolved. Because of the central role of acetyl-CoA in many metabolic pathways, it is likely that more than one pathway could contribute to maintaining the acetyl-CoA pool (Ohlrogge and Browse, 20 1995).

One school of thought takes the view that carbon for fatty acid synthesis is derived directly from the products of photosynthesis. In this scenario, 3-phosphoglycerate (3-PGA) would give rise to pyruvate, which would be converted to acetyl-CoA by pyruvate dehydrogenase in plastids 25 (Liedvogel, 1986). This hypothesis has many appealing aspects, but also several unaddressed questions: ( 1) fatty acid synthesis occurs in photosynthetic (chloroplasts) and non-photosynthetic plastids (in root, developing embryo cotyledons, endosperm leucoplasts); (2) some plastids may lack 3-phosphoglycerate mutase (Kleinig and Liedvogel, 1980), an 30 essential enzyme for converting 3-PGA, the immediate product of CO<sub>2</sub>

fixation, to pyruvate. (3) Acetate is the preferred substrate for fatty acid synthesis using isolated intact plastids, and there is evidence that a multienzyme system including acetyl-CoA synthetase and acetyl-CoA carboxylase, exists in plastids, which channels acetate into lipids (Roughan 5 and Ohlrogge, 1996). It is almost certain that at least some of the acetyl-CoA in plastids is formed by plastidic pyruvate dehydrogenase, using pyruvate imported from the cytosol or produced locally by plastidial glycolysis.

A further possibility, especially in non-photosynthetic tissues (e.g., 10 roots and developing embryos), is that acetyl-CoA, generated in the mitochondria, is an alternate means to provide acetate moieties for fatty acid synthesis (Ohlrogge and Browse, 1995). Mitochondrially-generated acetyl-CoA could be hydrolyzed to yield free acetate, which could move into the plastid for conversion to acetyl-CoA via plastidial acetyl-CoA synthetase, an 15 enzyme with 5- to 15-fold higher activity than the *in vivo* rate of fatty acid synthesis (Roughan and Ohlrogge, 1994). Alternatively, the mitochondrial acetyl-CoA could be converted to acetylcarnitine and transported directly into the plastid. Hence, in theory, the mitochondrial pyruvate dehydrogenase complex has an important role to play in fatty acid biosynthesis (see Fig. 1 of 20 the accompanying drawings). The proof of this hypothesis has been hindered by the difficulties of directly measuring the existence of acetate in the cytosol.

The mitochondrial PDC (mtPDC) is a tightly regulated multiple subunit complex. As mentioned previously, one of the key regulatory components of 25 this complex is PDH\_kinase (PDHK). PDHK functions as a negative regulator by inactivating PDH via phosphorylation. By modulating the PDCK, the activity of PDC can be genetically engineered.

Various attempts have been made to increase or channel additional carbon towards fatty acid biosynthesis. Targets have included genetically 30 modifying acetyl-CoA carboxylase and pyruvate kinase gene expression

through over-expression and antisense mRNA techniques with limited or no success.

However, there are many examples of successful modifications to plant metabolism that have been achieved by genetic engineering to transfer 5 new genes or to alter the expression of existing genes, in plants. It is now routinely possible to introduce genes into many plant species of agronomic significance to improve crop performance (e.g. seed oil or tuber starch content/composition; meal improvement; herbicide, disease or insect resistance; heavy metal tolerance etc.) (Somerville, 1993; Kishore and 10 Somerville, 1993; MacKenzie and Jain, 1997).

For example, increases in the proportions of some strategic fatty acids and in the quantities of seed oil have been achieved by the introduction of various fatty acid biosynthesis and acyltransferase genes in oilseed crops. These include the following demonstrations: Expression of an anti-sense 15 construct to the stearoyl-ACP  $\Delta 9$  desaturase in *Brassicaceae* led to an increase in the stearic acid content (Knutzon et al., 1992). Expression of a medium chain fatty acyl-ACP thioesterase from California Bay, in *Brassicaceae* was demonstrated to increase the lauric acid (12:0) content (Voelker et al., 1992; 1996). Expression of a *Jojoba*  $\beta$  keto-acyl-CoA 20 synthase in low erucic acid *Brassicaceae* led to an increase the level of erucic acid (22:1); the effect following expression in high erucic acid cultivars was negligible (Lassner et al., 1996). Increased proportions of oleic acid in *Brassica napus* and in soybean have been achieved by silencing the microsomal FAD2 ( $\Delta 12$ ) desaturase (Hitz et al., 1995; Kinney, 1995 ; 1997). 25 Transformation of *Arabidopsis thaliana* and rapeseed (*B. napus*) with a yeast *sn-2* acyltransferase resulted in seed oils with increased proportions of 22:1 and other very long-chain fatty acids and significant increases in seed oil content (Zou et al., 1997).

Starch deposition has also been altered by genetic engineering. By 30 expression of a mutant *E. coli glgC16* gene encoding an ADP glucose

pyrophosphorylase in potato tubers, an increase in starch accumulation was achieved (Stark et al., 1992).

However, because a PDHK gene has not heretofore been cloned from any plant, until now, no genetic modifications have addressed the possibility of altering carbon flux, increasing fatty acid synthesis, oil content or seed size, altering flowering and/or generation time, vegetative growth, or plant respiration/productivity by modulating plant mitochondrial PDH activity.

#### DISCLOSURE OF INVENTION

10

An object of the invention is to identify, isolate and characterize a pyruvate dehydrogenase kinase (PDHK) (gene and cDNA) sequence from *Arabidopsis* and to utilize this sequence in the genetic manipulation of plants.

Another object of the invention is to provide a vector containing the full-length PDHK sequence or a significant portion of the PDHK sequence from *Arabidopsis*, in an anti-sense orientation under control of either a constitutive or a seed-specific promoter, for re-introducing into *Arabidopsis* or for introducing into other plants.

Another object of the invention is to provide a method to construct a vector containing the full-length PDHK sequence or a significant portion of the PDHK sequence from *Arabidopsis*, in a sense orientation under control of either a constitutive or a seed-specific promoter, for re-introducing into *Arabidopsis* or for introducing into other plants.

Another object of the invention is to provide a method of modifying *Arabidopsis* and other plants to change their seed oil content.

Another object of the invention is to provide a method of modifying *Arabidopsis* and other plants to change their average seed weight or size.

Another object of the invention is to provide a method of modifying *Arabidopsis* and other plants to change their respiration rate during development.



Another object of the invention is to provide a method of modifying *Arabidopsis* and other plants to change their vegetative growth characteristics.

Another object of the invention is to provide a method of modifying  
5 *Arabidopsis* and other plants to change their flowering time or patterns of generative growth.

Yet another object of the invention is to provide a method of modifying *Arabidopsis* and other plants to change the period required to reach seed maturity.

10 According to one aspect of the present invention, there is provided isolated and purified deoxyribonucleic acid (DNA) of SEQ ID NO:1 (pYA5; ATCC No 209562).

According to yet another object of the invention, there is provided a vector containing SEQ ID NO:1 or a part thereof, for introduction of the gene,  
15 in an anti-sense orientation (e.g. pAsYA5; ATCC No 209561) into a plant cell, and a method for preparing a vector containing SEQ ID NO:1 or a part thereof, for introduction of the gene in a sense orientation, into a plant cell.

The invention also relates to transgenic plants and plant seeds having a genome containing an introduced DNA sequence of SEQ ID NO:1 and a  
20 method of producing such plants and plant seeds.

The invention also relates to substantially homologous DNA sequences from plants with deduced amino acid sequences of 25% or greater identity, and 50% or greater similarity, isolated and/or characterized by known methods using the sequence information of SEQ ID NO:1, as will  
25 be appreciated by persons skilled in the art, and to parts of reduced length that are still able to function as inhibitors of gene expression by use in an anti-sense or co-suppression (Transwitch; Jorgensen and Napoli 1994) application. It will be appreciated by persons skilled in the art that small changes in the identities of nucleotides in a specific gene sequence may  
30 result in reduced or enhanced effectiveness of the genes and that, in some

applications (e.g. anti-sense or co-suppression), partial sequences often work as effectively as full length versions. The ways in which the gene sequence can be varied or shortened are well known to persons skilled in the art, as are ways of testing the effectiveness of the altered genes. All 5 such variations of the genes are therefore claimed as part of the present invention.

Stated more generally, the present invention relates to the isolation, purification and characterization of a mitochondrial pyruvate dehydrogenase kinase (PDHK) gene from the *Brassicaceae* (specifically *Arabidopsis* 10 *thaliana*) and demonstrates its utility in regulating fatty acid synthesis, seed oil content, seed size/weight, flowering time, vegetative growth, respiration rate and generation time. Until now, no concrete data is available on the gene structure of plant PDC regulatory subunits (PDCK and PDC phosphatase).

15 The PDHK gene was cloned and characterized in the course of experiments designed to complement an *E. coli* mutant, JC201 (Coleman, 1990) with a plant (*A. thaliana*) cDNA library. By expressing the cDNA as a fusion protein in *E. coli*, its function was established as a PDHK in a protein kinase assay where it specifically phosphorylated the mammalian PDH 20 E1 $\alpha$ /E1 $\beta$  subunits (the specific substrates of PDHK). The *A. thaliana* PDHK structure is significantly homologous to its mammalian counterpart, particularly among the functional domains.

The PDHK of the invention is useful in manipulating PDH activity, and the respiration rate in plants. For example, by transforming plants with a 25 construct containing the partial PDHK gene in an antisense or in a sense orientation, under the control of either constitutive or tissue-specific promoters, the expression of mitochondrial PDHK can be silenced to some degree by anti-sense or co-suppression (Transwitch) phenomena (De Lange et al., 1995; Mol et al., 1990; Jorgensen and Napoli, 1994; Kinney, 1995), 30 respectively. This can result in increased mitochondrial PDH activity, and

hence an increased production or availability of mitochondrially-generated acetyl-CoA, or an increased respiration rate.

Alternatively, by over-expressing the full-length PDHK gene selectively in a tissue-specific manner, the activity of mitochondrial PDH may be negatively regulated, resulting in decreased respiratory rates in tissues, such as leaves or tubers, to decrease maintenance respiration and thereby increase the accumulation of biomass.

Some of the manipulations and deliverables which are possible using the PDHK gene or a part thereof, include, but are not limited to, the following: seeds with increased or decreased fatty acid and oil content; plants exhibiting early or delayed flowering times (measured in terms of days after planting or sowing seed); plants with increased or decreased vegetative growth (biomass); plants with root systems better able to withstand low soil temperatures or frost; plants with tissues exhibiting higher or lower rates of respiration; plants exhibiting an enhanced capacity to accumulate storage compounds in other storage organs (e.g. tubers); plants exhibiting an enhanced capacity to accumulate biopolymers which rely on acetyl moieties as precursors, such as polyhydroxyalkanoic acids or polyhydroxybutyric acids (Padgett et al., 1997).

20

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the central role played by acetyl-CoA in mitochondrial respiration and plastidial fatty acid biosynthesis. The pyruvate dehydrogenase complex (PDC) oxidatively decarboxylates pyruvate to yield acetyl-CoA. Plants are unique in that they have both mitochondrial and plastidial isoforms of the PDC. The mitochondrial pyruvate dehydrogenase complex plays a key role in the regulation of acetyl-CoA generation and availability of acetyl moieties for various catabolic and anabolic reactions in plant cells. The mitochondrial PDC is negatively regulated by

phosphorylation of the E1 $\alpha$  subunit by pyruvate dehydrogenase kinase (PDCK= PDHK), and positively regulated by dephosphorylation of the PDC by pyruvate dehydrogenase phosphatase (PDCP). Mitochondrially-generated acetyl moieties can find their way into the respiratory tricarboxylic acid (TCA) cycle, but also into the plastid compartment where ultimately, acetate units are used by the enzymes of the fatty acid synthesis (FAS) pathway to synthesize fatty acids. These are eventually incorporated into membrane and also storage glycerolipids. Other abbreviations: PDC, pyruvate dehydrogenase complex; OAA, oxaloacetate; ACS, acetyl-CoA synthetase; ACH, acetyl-CoA hydrolase; DHAP, dihydroxyacetone phosphate.

Figure 2 shows the nucleotide sequence [SEQ ID NO:1] and the deduced amino acid sequence [SEQ ID NO:2] of the *Arabidopsis thaliana* PDH kinase (PDHK) cDNA (clone YA5).

Figure 3 shows the amino acid sequence alignment of the *Arabidopsis* PDH kinase (Ya5p) [SEQ ID NO:2] with other mammalian mitochondrial ketoacid dehydrogenase kinases: **Pdhk I**, porcine PDH kinase subunit I [SEQ ID NO:3]; **Pdhk II**, porcine PDH kinase subunit II [SEQ ID NO:4]; and **Bckdhk**, porcine branched chain  $\alpha$ -ketoacid dehydrogenase kinase [SEQ ID NO: 5]. Dots indicate gaps. Identical amino acid residues are highlighted in bold upper case type.

Figure 4 shows the predicted helical wheel structure (angle = 100°) of the 24 amino acid residues at the N-terminus of the YA5 (pyruvate dehydrogenase kinase, PDHK) protein. The N-terminal leader sequence of the YA5 protein corresponds well to most mitochondrial targeting sequences (Rosie and Schatz, 1988), consisting of a stretch of amino acids enriched in hydrophobic residues and opposing positively-charged residues. The key hydrophobic

(V<sub>3</sub>, F<sub>10</sub>, L<sub>14</sub>, V<sub>18</sub>, and W<sub>21</sub>) and positively-charged (K<sub>5</sub>, K<sub>12</sub> and H<sub>19</sub>) residues are found on opposing sides of the helical wheel motif in this mitochondrial targeting sequence, and are highlighted by • on the residue itself, and by ? next to the residue number.

5

Figure 5 shows the results of a DNA gel blot analysis (Southern, 1975) of the *Arabidopsis thaliana* YA5 (PDHK) gene. Genomic DNA was digested with PstI + XbaI (lane 1), XbaI (lane 2), PstI (lane 3), PvuII + SpeI (lane 4), SpeI (lane 5), and PvuII (lane 6). None of these enzymes has an internal 10 restriction site on the YA5 (PDHK) cDNA. The digested DNA was hybridized with <sup>32</sup>P-labeled YA5 cDNA (≈1.5 Kb) under high stringency conditions. All digests show only one hybridizing fragment suggesting that the PDHK gene most likely represents a single-copy gene in *Arabidopsis thaliana*.

15 Figure 6 depicts an RNA (northern) gel blot analysis of YA5 (PDHK) mRNA abundance/tissue distribution in *A. thaliana*. RNA was extracted from flowers (F), vegetative tissue (seedling leaves (L)), young developing siliques (YS) and maturing siliques (MS). The analysis shows that in all tissues, an RNA-hybridizing band of about 1.5 Kb was observed, but the abundance of the 20 YA5 mRNA varied considerably from tissue to tissue. Young seedling leaves (L) showed the highest level of YA5 expression, while significant, but lower, levels of expression were observed in developing siliques (seeds).

Figure 7a, 7b, 7c and 7d show the results of experiments in which the YA5 25 PDHK cDNA was expressed as a fusion protein in *E. coli*. and tests conducted to confirm its function as a PDH Kinase. The YA5 cDNA was expressed as a fusion protein in *E. coli* as shown in Fig. 7a. Analysis by SDS-Polyacrylamide Gel Electrophoresis (SDS-PAGE) revealed that lysate of *E. coli* transformed with the *A. thaliana* PDHK (YA5) has a very strongly- 30 induced fusion protein of M<sub>r</sub> ≈ 45 kDa (?), which is the predicted mass of the

*A. thaliana* PDHK gene fusion product (42 kD + 3 kD His TAG). Fig. 7b shows the mammalian E1 $\alpha$ /E1 $\beta$  PDH subunit complex (obtained courtesy of Dr. M. Patel at the University of Buffalo). The proteins have been co-expressed in *E. coli*, to provide a substrate to test the capacity of PDHK to phosphorylate the E1 subunit of the PDH complex. Fig. 7c and 7d are autoradiograms of radioactive incorporation of  $^{32}\text{P}$  (from  $\gamma\text{-}^{32}\text{P}\text{-ATP}$ ) into the E1 subunit of the E1 $\alpha$ /E1 $\beta$  PDH complex. The left hand panels of 7c and 7d show the time-dependent (incubation times of 2, 5, 10, 15, or 20 min) in vitro phosphorylation of the E1 $\alpha$ /E1 $\beta$  PDH complex by the action of the plant PDHK (product of clone YA5 expressed in *E. coli*), confirming its function as a pyruvate dehydrogenase kinase the first cloned from plants. In Fig. 7c the control reaction (right hand panel) contains YA5 lysate + control *E. coli* lysate without E1 $\alpha$ /E1 $\beta$  substrate. There is no evidence of phosphorylation of the E1 $\alpha$ /E1 $\beta$  complex. In Fig. 7d the control reaction (right hand panel) contains control *E. coli* lysate (without YA5 insert) + the E1 $\alpha$ /E1 $\beta$  substrate. Again, there is no evidence of phosphorylation of the E1 $\alpha$ /E1 $\beta$  complex.

Figure 8 shows the mitochondrial pyruvate dehydrogenase (PDC) activity in leaves from *A. thaliana* non-transformed wild-type (n-WT) plants and  $T_2$  transgenic plants containing constitutively expressed anti-sense pyruvate dehydrogenase kinase (PDHK), designated as YA5 lines. Mitochondria isolated from leaves of *A. thaliana* YA5 transgenic lines containing a constitutively-expressed anti-sense PDHK construct, have elevated activity of PDC compared to mitochondria isolated from leaves of non-transformed control plants.

Figure 9 shows the mitochondrial citrate synthase activity in leaves from *A. thaliana* non-transformed wild-type (n-WT) plants and  $T_2$  transgenic plants containing constitutively expressed anti-sense pyruvate dehydrogenase kinase (PDHK), designated as YA5 lines. Mitochondria isolated from leaves

of *A. thaliana* transgenic lines transformed with a constitutively-expressed anti-sense PDHK construct also have elevated activities of citrate synthase, in addition to elevated PDC, compared to mitochondria isolated from leaves of non-transformed control plants.

5

Figure 10 shows the oil content (expressed as  $\mu\text{g}$  total fatty acids per 100 seeds) in seeds isolated from *A. thaliana* non-transformed controls (nt-WT Con), and  $T_2$  seeds of pBI121 plasmid only control (pBI121 Con) transgenics, and anti-sense pyruvate dehydrogenase kinase (PDHK) transgenics, designated as YA5 lines. The *A. thaliana* YA5 seed lines transformed with a constitutively-expressed anti-sense PDHK construct have elevated fatty acid and oil content compared to seeds from non-transformed control plants or transformants containing only the selectable marker gene (transformed with pBI121), but without anti-sense PDHK.

15

Figure 11 shows the time (expressed in days after planting) to reach the flower initiation (generative) phase in *A. thaliana* non-transformed controls (nt-WT), and the  $T_2$  generation of pBI121 plasmid only control (pBI121) transgenics, and anti-sense pyruvate dehydrogenase kinase (PDHK) transgenics, designated as YA5 lines. The time to reach the generative (floral initiation) stage is reduced in *A. thaliana* YA5 lines transformed with a constitutively-expressed anti-sense PDHK construct, compared to non-transformed controls or transformants containing only the selectable marker gene (transformed with pBI121), but without anti-sense PDHK.

25

Figure 12 shows vegetative shoot tissue dry weights at 31 days after planting in *A. thaliana* non-transformed controls (WT), and the  $T_2$  generation of pBI121 plasmid only control (pBI121) transgenics, and anti-sense pyruvate dehydrogenase kinase (PDHK) transgenics, designated as YA5 lines. Shoot tissue growth is reduced in *A. thaliana* YA5 lines transformed with a

constitutively-expressed anti-sense PDHK construct, compared to non-transformed controls or transformants containing only the selectable marker gene (transformed with pBI121), but without anti-sense PDHK.

5 Figure 13 shows the average number of rosette leaves present upon entering the generative phase, in *A. thaliana* non-transformed control (WT) plants, and the T<sub>2</sub> generation of pBI121 plasmid only control (pBI121) transgenics, and anti-sense pyruvate dehydrogenase kinase (PDHK) transgenics, designated as YA5 lines. The average number of rosette  
10 leaves per plant is reduced in *A. thaliana* YA5 lines transformed with a constitutively-expressed anti-sense PDHK construct, compared to non-transformed controls or transformants containing only the selectable marker gene (transformed with pBI121), but without anti-sense PDHK. The  
15 perturbed vegetative growth phase in the anti-sense PDHK transgenics (see also Fig. 12 ) correlates well with the earlier flowering phenotype (see also Fig. 11).

#### BEST MODES FOR CARRYING OUT THE INVENTION

20

The best modes for carrying out the invention are apparent from the following description of the results of tests and experiments that have been carried out by the inventors.

The inventors chose to use the well-accepted model plant system  
25 *Arabidopsis thaliana* for the cloning of PDHK, as a host system for genetic engineering to alter PDHK expression, and to study the effects of altering PDHK expression on various plant developmental and metabolic processes.

This is because over the past several years, *Arabidopsis thaliana*, a typical flowering plant, has gained increasing popularity as a model system  
30 for the study of plant biology. As a result of the ease with which this plant



lends itself to work in both classical and molecular genetics, *Arabidopsis* has come to be widely used as a model organism in plant molecular genetics, development, physiology and biochemistry (Meyerowitz and Chang, 1985; Meyerowitz, 1987; Goodman et al., 1995). This model dicotyledonous plant is also closely related to *Brassica* crop species and it is increasingly apparent that information concerning the genetic control of basic biological processes in *Arabidopsis* will be transferable to other species (Lagercrantz et al., 1996).

Indeed, there are numerous examples wherein studies of the molecular biology and biochemistry of a particular metabolic pathway or developmental process and the possibility of genetically engineering a plant to bring about changes to said metabolic pathway or process, has first been tested in the model plant *Arabidopsis*, and then shown to yield similar phenotypes in other plants, particularly crop plants.

For example, the extra- plastidial membrane associated oleate (18:1)  $\Delta 12$  ( $\omega$ -6) desaturase gene, FAD2, was originally studied and eventually cloned from *Arabidopsis thaliana*, by identifying the lesion found in an *A. thaliana* mutant defective in desaturating oleate to produce linoleate (18:2) on the phosphatidylcholine backbone. This resulted in a high oleic acid phenotype in the *A. thaliana* seed oil (Okuley et al., 1994). Genetic engineering of both soybean (*Glycine max.*) and canola *B. napus* to silence the indigenous FAD2 gene(s) in a seed-specific manner by anti-sense or co-suppression approaches, resulted in similar high oleic acid seed oil phenotypes (Kinney, 1995; 1997).

Transgenic expression of a yeast *sn*-2 acyltransferase (*SLC1-1*) gene to achieve enhanced seed oil and very long-chain fatty acid content was first performed in *Arabidopsis* and later shown to yield similar phenotypes in transgenic rapeseed (*B. napus*) experiments (Zou et al., 1997). *Arabidopsis thaliana* has repeatedly shown itself to be a useful model system for metabolic engineering of metabolic pathways (e.g. lipid biosynthesis,

photosynthesis) or processes (organogenesis, reproductive development etc.) common to all higher plants.

In the area of secondary metabolism/signal transduction, an anthocyanin pathway-specific transcriptional activator from the monocot 5 maize designated as *R* (the *myc* transcription factor involved in activation of biosynthetic genes for anthocyanin production in the aleurone cells of maize kernels), was expressed in the dicot *Arabidopsis*, causing augmented anthocyanin pigmentation in the inflorescences. Subsequent expression in another dicot, tobacco (*Nicotiana tabacum*), resulted in similar floral 10 pigmentation changes (Lloyd et al., 1992). These experiments demonstrate that whole pathways common to all flowering plants can be co-ordinately controlled through the introduction of transcriptional regulators, and that the mechanisms are common to diverse plant species.

In the context of the current invention, all plant cells undergo 15 mitochondrial respiration and this ubiquitous process is affected by the activity of the PDC and its regulators PDCK and PDCP as explained previously. Thus, many of the effects observed following genetic engineering to modulate PDCK expression in *Arabidopsis* can be expected to result in similar phenotypes when carried out in all other plants.

20 There are a number of ways by which genes and gene constructs can be introduced into plants, and a combination of plant transformation and tissue culture techniques have been successfully integrated into effective strategies for creating transgenic crop plants. These methods, which can be used in the present invention, have been extensively reviewed elsewhere 25 (Potrykus, 1991; Vasil, 1994; Walden and Wingender, 1995; Songstad et al., 1995), and are well known to persons skilled in the art. For example, one skilled in the art will certainly be aware that, in addition to *Agrobacterium*-mediated transformation of *Arabidopsis* by vacuum infiltration (Bechtold et al., 1993) or wound inoculation (Katavic et al., 1994), it is equally possible to 30 transform other plant and crop species, using *Agrobacterium* Ti-plasmid-

mediated transformation (e.g. hypocotyl (DeBlock et al., 1989) or cotyledonary petiole (Moloney et al., 1989) wound infection), particle bombardment/biolistic methods (Sanford et al., 1987; Nehra et al., 1994; Becker et al., 1994) or polyethylene glycol-assisted protoplast transformation 5 (Rhodes et al., 1988; Shimamoto et al., 1989) methods.

As will also be apparent to persons skilled in the art, and as extensively reviewed elsewhere (Meyer, 1995; Datla et al., 1997), it is possible to utilize plant promoters to direct any intended up- or down-regulation of transgene expression using constitutive promoters (e.g. those 10 based on CaMV35S), or by using promoters which can target gene expression to particular cells, tissues (e.g. napin promoter for expression of transgenes in developing seed cotyledons), organs (e.g. roots), to a particular developmental stage, or in response to a particular external stimulus (e.g. heat shock).

15 Particularly preferred plants for modification according to the present invention include borage (*Borago* spp.), Canola, castor (*Ricinus communis*), cocoa bean (*Theobroma cacao*), corn (*Zea mays*), cotton (*Gossypium* spp), *Crambe* spp., *Cuphea* spp., flax (*Linum* spp.), *Lesquerella* and *Limnanthes* spp., Linola, nasturtium (*Tropaeolum* spp.), *Oenothera* spp., olive (*Olea* 20 spp.), palm (*Elaeis* spp.), peanut (*Arachis* spp.), rapeseed, safflower (*Carthamus* spp.), soybean (*Glycine* and *Soja* spp.), sunflower (*Helianthus* spp.), tobacco (*Nicotiana* spp.), *Vernonia* spp., wheat (*Triticum* spp.), barley (*Hordeum* spp.), rice (*Oryza* spp.), oat (*Avena* spp.) sorghum (*Sorghum* spp.), rye (*Secale* spp.) or other members of the *Gramineae*.

## 25 RESULTS

### *cDNA cloning and sequence analysis of clone YA5 (plant PDHK).*

A plant PDHK cDNA sequence designated YA5 was identified and 30 cloned during experiments designed to complement an *E. coli* mutant JC201 (Coleman, 1990) with an *Arabidopsis thaliana* cDNA library. The *E. coli*

mutant JC201 has been reported to be a mutant deficient in *lyso*-phosphatidic acid acyltransferase (LPAT; EC 2.3.1.51) activity, and possesses a temperature-sensitive growth phenotype (Coleman, 1990). Plasmids generated from an *A. thaliana*  $\lambda$ -YES expression library (Elledge et al., 1991) were used to transform *E. coli* mutant JC201. A restrictive temperature condition (44°C) was applied to select surviving colonies. cDNAs were isolated from temperature-insensitive transformants. Clone YA5 was found to be able to complement or rescue the temperature-sensitivity of JC201, but no elevated LPAT activity could be detected in 10 lysates of the transformant. Thus, the mechanism underlying the ability to complement the temperature sensitivity of JC201 remains unclear. However, several other complementing clones have also been found to rescue the temperature-sensitive phenotype of JC201, indicating that temperature complementation can occur upon transformation with cDNAs having 15 functions unrelated to LPAT (Taylor et al., 1992a; Zou and Taylor, 1994).

The YA5 cDNA was sequenced from both strands on an Applied Biosystems Model 373A DNA Sequencing System using the *Taq* DyeDeoxy™ Terminator Cycle Sequencing Kit (Applied Biosystems, Inc.). The nucleotide sequence of the 1.457 kb YA5 cDNA (pYA5; ATCC 209562) 20 [SEQ ID NO:1] and its deduced amino acid sequence [SEQ ID NO:2] are shown in Figure 2. A sample of the YA5 cDNA (pYA5) was deposited on December 18, 1997 at the American Type Culture Collection (ATCC) of 12301 Parklawn Drive, Rockville, Maryland 20852, USA, under accession number ATCC 209562. The sequence revealed a 103 nucleotide 5' 25 untranslated region, and a 235 nucleotide 3' untranslated region followed by a poly A tail. YA5 has an open reading frame of 1098 base pairs encoding a polypeptide of 366 amino acids, with a calculated molecular weight of 41.37 kDa. The sequences around the initiation codon AUG are in good agreement with the consensus sequences derived from other plant species 30 (Lutcke et al., 1987). There is an in-frame stop codon upstream of the start

codon, indicating that YA5 is a full-length cDNA. The calculated isoelectric point of the YA5 protein is 6.68 and its net charge at pH 7.0 is calculated to be -1.48.

### 5 Amino Acid Sequence Alignment

As shown in Figure 3, comparisons of the deduced amino acid sequence [SEQ ID NO:2] of the YA5 protein (Ya5p) to the NCBI data bank revealed a high degree of homology with mammalian mitochondrial kinases responsible for phosphorylation and inactivation of  $\alpha$ -ketoacid dehydrogenase complexes (Harris et al., 1992), including the pyruvate dehydrogenase complex (PDC), the  $\alpha$ -ketoglutarate dehydrogenase complex (KGDC) and the branched-chain  $\alpha$ -ketoacid dehydrogenase complex (BCKDHC). These mammalian complexes are located in the mitochondrial matrix space (Damuni et al, 1984) and are similar in both structure and function (Nobukuni et al., 1990). cDNAs encoding the mammalian pyruvate dehydrogenase kinase (PDHK) and the branched-chain  $\alpha$ -ketoacid dehydrogenase kinase (BCKDHK) have been cloned and the amino acid sequences of these protein kinases are highly homologous to each other (Popov et al., 1992; 1993; 1994).

The YA5 protein (Ya5p) is 28.6% identical and 83.7% similar to PDKI (Popov et al., 1993) and 32.3% identical and 88.4% similar to PDKII (Popov et al., 1994), both subunits of the porcine PDH kinase. Ya5p is also 28.8% identical and 84.1% similar to BCKDHK (Popov et al., 1992). The sequence similarity extends over the entire sequence, but sequence differences and alignment gaps occur throughout, particularly towards the amino and carboxyl termini.

SEQ ID NO:1 of the current invention and mammalian PDHK and BCKDHK do not exhibit significant homology with known serine/threonine protein kinases. Rather, a much higher degree of sequence homology was found with members of the prokaryotic protein histidine kinase family. As

shown in Figure 3, the most homologous regions fall into the conserved motifs defining histidine kinase functional domains. Members of the protein histidine kinase family have five regions that are highly conserved (Parkinson and Kofoed, 1992). All five motifs are easily identifiable in the YA5 deduced amino acid sequence, with the same order and spacing conserved in bacterial proteins. At the C-terminus, the catalytic domain (Block V) with a glycine-rich loop of Gly<sup>320</sup>-X-Gly<sup>322</sup>-X-Gly<sup>324</sup> [SEQ ID NO:7] as well as the sequence surrounding it, is the longest stretch of amino acids that exhibit high identity. Block III with the consensus sequence of Asp<sup>278</sup>-X-Gly<sup>280</sup>-X-Gly<sup>282</sup> [SEQ ID NO:8] characteristic of adenosine triphosphate (ATP)-binding proteins, and Block IV with an invariant Phe<sup>292</sup>, are located at the positions defined as the central core of the catalytic domain. A highly-conserved region defined as Block II (Glu<sup>238</sup>-Leu-X-Lys-Asn<sup>242</sup>-X-X-Arg-Ala<sup>246</sup>) [SEQ ID NO: 9] of the catalytic domain is also found at the proper proximity to the N-terminus. The histidine residue (His<sup>121</sup>) conserved among YA5, PDKI and PDKII would probably represent Block I, that is proposed to be involved in autophosphorylation.

The N-terminal leader sequence of the YA5 protein corresponds well to most mitochondrial targeting sequences (Rosie and Schatz, 1988), consisting of a stretch of amino acids enriched in hydrophobic and positively-charged residues, with a predicted helical wheel structure (angle = 100°) (Figure 4). The key hydrophobic (V<sub>3</sub>, F<sub>10</sub>, L<sub>14</sub>, V<sub>18</sub>, and W<sub>21</sub>) and positively-charged (K<sub>5</sub>, K<sub>12</sub> and H<sub>19</sub>) residues are found on opposing sides of the helical wheel motif in this mitochondrial targeting sequence. The YA5 protein lacks obvious targeting motifs typically found in proteins targeted to the peroxisomes (e.g. extreme C-terminus non-cleaved Ser-Lys-Leu (SKL) peroxisomal targeting sequence motif; Mullen et al., 1997).

### *Genomic organization and expression of the YA5 gene in A. thaliana*

Genomic DNA was digested with [ XbaI + PstI], XbaI, PstI, [PvuII + SpeI], SpeI, and PvuII (there are no internal restriction sites on the YA5 cDNA for these enzymes). The digested DNA was then subjected to a DNA gel blot (Southern, 1975) and hybridized with the <sup>32</sup>P-labeled YA5 cDNA under high stringency hybridization conditions. As shown in Figure 5, all of the digest reactions yielded only one hybridizing fragment (gel band), indicating that the YA5 gene most likely is present as a single copy in the *Arabidopsis* genome.

To determine the relative abundance and tissue distribution of YA5 gene transcript, an RNA gel blot (northern blot) hybridization analysis, shown in Figure 6, was performed on RNA extracted from *A. thaliana* seedlings, inflorescences (flowers), young siliques and maturing siliques. In all tissues, an RNA-hybridizing band of about 1.5 kb was observed, but the abundance of the YA5 mRNA varied considerably from tissue to tissue. Young seedlings showed the highest level of YA5 expression, while significant, but lower levels of expression were observed in developing siliques (seeds).

### *Expression of YA5 in E. coli and Confirmation of its Function as a PDH Kinase*

The YA5 full-length cDNA (YA5F) was cloned in pBluescript SK in a 5' to 3' orientation of T7-T3. A primer encompassing the putative translational initiation site Ompdk (AGGAGAGACTCGAGGCTTATGGCAGTGAAG) [SEQ ID NO:6] was synthesized to include a XhoI restriction site. Primer Ompdk and T3 primer were used in a PCR reaction to amplify the YA5 encoding region from YA5F. The resulting PCR fragment was digested with HindIII (HindIII site is present 3' to the stop codon) and XhoI and cloned into pTrcHisB vector (Clontech) to generate construct pZTa5. SDS-PAGE analysis with lysates from IPTG induced *E. coli* containing pZTa5 and

pTrcHisB control vector (Figure 7a), lanes 1 ("YA5") and 2 ("Control"), respectively), revealed that the pZTa5 transformant (Fig. 7a, lane 1 "YA5") exhibited a very strongly-induced fusion protein of  $M_r \approx 45$  kD, which is the predicted mass of the *A. thaliana* PDHK gene fusion product ( $\approx 42$  kD + 3 kD His TAG).

Protein kinase activity of the *E. coli* expressed YA5 protein was assayed essentially as described by Liu et al., (1995). The protein phosphorylation substrates, human E1 $\alpha$  and E1 $\beta$  co-expressed in, and purified from, *E. coli* M15, were obtained from Dr. M.S. Patel of the Department of Biochemistry, School of Medicine and Biomedical Sciences, State University of New York at Buffalo, Buffalo, New York (Figure 7b). This co-expressed E1 $\alpha$  and E1 $\beta$  system has been used extensively in the study of the regulation of PDC E<sub>1</sub> phosphorylation in mammalian systems (Korotchkina and Patel, 1995). For phosphorylation experiments (shown in Fig. 7c and 7d), 20-25  $\mu$ g of E1 $\alpha$ /E1 $\beta$  was combined with about 10  $\mu$ g of YA5-accumulating *E. coli* cytosol protein in a final volume of 100  $\mu$ L containing 20 mM potassium phosphate, pH 7.0, 1mM magnesium chloride, 2 mM dithiothreitol, 0.1 mM EDTA and 200  $\mu$ M cold ATP. The mixture was pre-incubated at room temperature for 5 minutes, and then 5 $\mu$ Ci  $^{32}$ P- $\gamma$ -ATP was added to start the assay. After 2, 5, 10, 15 and 20 minutes, 20- $\mu$ L aliquots were withdrawn and the reaction was stopped with 20  $\mu$ L of SDS-denaturing mixture. Samples were separated on 10% SDS-PAGE and autoradiographed.

Figures 7c and 7d are autoradiograms of radioactive incorporation of  $^{32}$ P (from  $\gamma$ - $^{32}$ P-ATP) into the E1 subunit of the E1 $\alpha$ /E1 $\beta$  complex. The left hand panels of 6(c) and 6(d) show the time-dependent (incubation times of 2, 5, 10, 15, or 20 min) *in vitro* phosphorylation of the E<sub>1</sub> $\alpha$ /E<sub>1</sub> $\beta$  complex substrate by the action of the plant PDHK (clone YA5) fusion protein, confirming its function as a pyruvate dehydrogenase kinase, the first cloned from plants. In Fig. 7c the control reaction (right hand panel) contains YA5



lysate + control *E. coli* lysate without PDH E1 $\alpha$ /E1 $\beta$  substrate. There is no evidence of phosphorylation of the E1 $\alpha$ /E1 $\beta$  substrate. In Fig. 7d the control reaction (right hand panel) contains control *E. coli* lysate (without YA5 insert) + the E1 $\alpha$ /E1 $\beta$  PDH complex substrate. There is no evidence of 5 phosphorylation of the E1 $\alpha$ /E1 $\beta$  complex in this control either.

*Synthesis of YA5 Plant Transformation Constructs:*

*Anti-sense YA5 (Anti-sense PDHK) Construct for Constitutive*

*10 Expression:*

The YA5 cDNA contains internal BamHI (nt 628) and NcoI (nt 1176) restriction sites. The BamHI and NcoI fragment was freed from YA5F and cloned into the respective sites in pBI524 (Datla et al., 1993), in an anti-sense orientation, and located between the tandem cauliflower mosaic virus 15 35S promoter and the nopaline synthase terminator. The YA5 anti-sense cassette was then cut out from pBI524 by HindIII and EcoRI, and cloned into the respective sites of vector pRD400 (Datla et al., 1992). The final binary anti-sense vector pAsYA5/pRD400 (a sample of which was deposited on December 18, 1997 under the terms of the Budapest Treaty at the American 20 Type Culture Collection (ATCC) of 12301 Parklawn Drive, Rockville, Maryland 20852, USA, under accession number ATCC 209561) was introduced into *Agrobacterium tumefaciens* strain GV3101 (bearing helper plasmid pMP90; Koncz and Schell, 1986) by electroporation.

25 *Anti-Sense and Partial Sense YA5 (PDHK) Constructs for Seed-Specific Expression:*

An 875 bp fragment of the YA5 cDNA was excised by a BamHI (Pharmacia) digestion and the fragment was ligated into plasmid pDH1, which contains the seed-specific napin promoter (pDH1 was kindly provided 30 by Dr. P.S. Covello, NRC/PBI). The cassette and the insert (in either sense

or anti-sense orientation) were excised by a partial digestion with HindIII and EcoRI, and the DNA fragments separated on agarose gels and purified using the GeneClean II Kit (Bio 101 Inc.). The fragments were then ligated into HindIII/EcoRI-digested pRD400. The final binary vectors pNAsYA5/pRD400 5 (anti-sense construct) or pNSYA5/pRD400 (partial sense construct), were introduced into *Agrobacterium tumefaciens* strain GV3101 (bearing helper plasmid pMP90; Koncz and Schell, 1986) by electroporation.

10 *Constitutive Expression of Anti-Sense YA5 (anti-sense PDHK) Gene in Arabidopsis thaliana:*

*Agrobacterium* containing the pAsYA5/pRD400 was used to transform *Arabidopsis* by vacuum infiltration (Bechtold et al., 1993). In addition, it will 15 be apparent to persons skilled in the art that transformation of *Arabidopsis* can also be achieved by wound inoculation (Katavic et al., 1994). Similarly, one skilled in the art will certainly be aware that transformation of other plant species is possible using *Agrobacterium* Ti-plasmid-mediated transformation (e.g. hypocotyl (DeBlock et al., 1989) or cotyledonary petiole (Moloney et al, 20 1989) wound infection), particle bombardment/biolistic methods (Sanford et al., 1987; Nehra et al., 1994; Becker et al., 1994) or polyethylene glycol-assisted protoplast transformation (Rhodes et al., 1988; Shimamoto et al., 1989) methods. Constructs may be driven by constitutive or tissue-specific (e.g. seed, root etc.) promoters, as will also be apparent to persons skilled in 25 the art.

As controls, plants were either non-transformed (nt), or transformed with pBI121 vector only (Jefferson et al., 1987; without antisense-PDHK insert but containing the kanamycin selection marker and the  $\beta$ -glucuronidase reporter gene). Control and transgenic plants were grown at

the same time under identical conditions in growth chambers as described by Katavic et al., 1995.

Results of DNA gel blot (Southern, 1975) analyses confirmed that all of the anti-sense PDHK transgenic lines (designated as YA5 lines 23, 31, 32, 5 52, 95, 104) have at least one insert per genome for the PDHK gene in an anti-sense orientation. As expected, the non-transformed wild type control and pBI121 (plasmid only) transgenic control have only one insert per genome, consistent with the original Southern analysis (see Figure 5).

10

*Analysis of Pyruvate Dehydrogenase (PDH) Activity in Mitochondria Isolated from A. thaliana Anti-Sense PDHK Transgenic Plants*

Shoot tissue was collected from *A. thaliana* transgenic plants 15 containing the anti-sense PDHK construct, and from non-transformed control plants, and intact mitochondria were isolated. Pyruvate dehydrogenase (PDH) activity was determined by the method of Reid et al., (1977). As shown in Figure 8, the PDH activity in mitochondria isolated from leaves of anti-sense PDHK transgenic plants was elevated by 20 to 350%, compared 20 to PDH activity in mitochondria isolated from non-transformed controls.

*Analysis of Kreb's Cycle Enzyme Activities in Mitochondria Isolated from A. thaliana Anti-Sense PDHK Transgenic Plants*

25 The activities of the Kreb's Cycle enzymes citrate synthase, fumarase and succinate dehydrogenase were all significantly elevated in mitochondria isolated from leaves of anti-sense PDHK transgenic plants, compared to the respective activities in mitochondria isolated from non-transformed wild-type (n-WT) controls. Citrate synthase activities were about 160-240% higher 30 (Figure 9), while fumarase activities were about 65-120% higher, and

succinate dehydrogenase activities were about 10-65% higher in the anti-sense PDHK transgenics, compared to the corresponding n-WT activities set at 100%. These results suggest that mitochondrial respiration is increased in the anti-sense PDHK transgenics due to an increased availability of acetyl-5 CoA generated by enhanced PDC activity (due to down regulation of expression of PDHK, a negative regulator of PDC).

*Analysis of Fatty Acid Composition and Content of Oils, and Average Seed Weights in T<sub>2</sub> Seeds from A. thaliana Anti-Sense PDHK and pBI121 Control*  
10 *Transgenic Plants*

Mature siliques and seeds were isolated from anti-sense PDHK transformants and from controls, either non-transformed or pBI121 transformants (without anti-sense PDHK, but with a kanamycin resistance 15 gene), and the respective oil contents, fatty acyl compositions of the seed oils, average seed weights and number of siliques per 15 cm segment of stem were determined.

As shown in Figure 10, overall oil content, expressed as µg total fatty acids/ 100 seeds, was significantly elevated in the anti-sense PDHK 20 transformants, by 8.5-26.5 %, and by 15.4-34.6%, compared to pBI121 transformants and non-transformed controls, respectively. This indicated that overall flux of acetyl moieties into seed storage lipids was enhanced by a greater contribution from mitochondrially-generated acetate. The latter was enabled by increased mitochondrial PDH activity, due to the down-regulation 25 of the negative regulator PDHK, in the antisense PDHK transformants.

Table 1 shows the oil content and average weight of seeds isolated from *A. thaliana* lines transformed with a constitutively-expressed anti-sense PDHK construct in comparison with seeds of plasmid only (pBI121) transformants and non-transformed controls. Both the amount of oil and the 30 average seed weight are higher in the anti-sense PDHK transformants.

Table 1

Mean seed oil content and seed weight in non-transformed controls and T<sub>2</sub> seed of pBI121 control and anti-sense YA5 (A/S PDHK) transgenic plants.

5

<i>A. thaliana</i> line	Seed Oil Content (mg oil /400 seeds)	Seed Weight (mg/ 400 seeds)
Non-transformed Control	3.13	7.40
pBI121 (plasmid only) Control	3.30	7.21
A/S YA5 31	3.89	9.02
A/S YA5 32	3.66	8.55
A/S YA5 52	3.75	8.61
A/S YA5 95	3.68	8.70
A/S YA5 104	3.87	8.83

The average number of siliques per 15 cm segment of bolted stem 10 was not significantly affected in the *A. thaliana* lines transformed with a constitutively-expressed anti-sense PDHK construct (designated as YA5 lines), in comparison with plasmid only (pBI121) transformants and non-transformed control plants. Following propagation of the T<sub>2</sub> generation of seed, non-transformed wild-type and pBI121 control transgenic *A. thaliana* 15 plants produced  $30 \pm 3$  siliques and  $31 \pm 4$  siliques, respectively, per 15 cm segment of bolted stem. The anti-sense PDHK transgenic YA5 lines 31, 32, 52, 95 and 104 produced  $26 \pm 3$ ,  $27 \pm 3$ ,  $27 \pm 3$ ,  $26 \pm 3$ , and  $24 \pm 3$  siliques per 15 cm segment of bolted stem, respectively.

The average number of  $T_3$  seeds per silique was also not significantly affected. For example, the pBI121 control transformant produced  $49.4 \pm 6.6$   $T_3$  seeds per silique, and the anti-sense PDHK line YA 95 produced  $50.1 \pm 8.5$   $T_3$  seeds per silique ( $n = 5-6$  mature siliques sampled from 4 individual transgenic plants of each line). This indicated that seed yield (harvest index) was not adversely affected in the anti-sense PDHK transformants.

Table 2 shows the fatty acyl composition of seed oils isolated from *A. thaliana* lines transformed with a constitutively-expressed anti-sense PDHK construct in comparison with oil from seeds of plasmid only (pBI121) transformants and non-transformed controls. The antisense PDHK construct affects a point very early in the fatty acid biosynthesis/lipid bioassembly pathway, i.e. it enables a greater availability of acetyl moieties for plastidial fatty acid biosynthesis. Thus, while total flux of carbon through the lipid pathway into storage lipids was enhanced in seeds of the anti-sense PDHK transgenic plants, the fatty acyl composition of the oils which accumulated was not markedly changed (Table 2).

Table 2  
 Acyl composition of oils from seed of non-transformed controls (n-WT Control) and T<sub>2</sub> seed of pBI121 control (pBI121 Control) and Anti-Sense PDHK (A/S YA5) transgenic plants.

Sample	16:0	18:0	18:1	18:2	18:3	20:0	20:1 c11+c13	20:2	22:0	22:1 c13+c15	22:2	24:0	24:1
n-WT Control	7.5 (0.2)*	3.5 (0.2)	14.9 (0.2)	27.0 (0.06)	20.1 (0.04)	1.9 (0.03)	20.4 (0.1)	2.2 (0.06)	0.4 (0.1)	1.9 (0.1)	0.1 (0.1)	0.1 (0.1)	0.1 (0.1)
pBI121 Control	7.7 (0.01)	3.4 (0.02)	15.5 (0.1)	27.3 (0.2)	19.8 (0.2)	1.8 (0.0)	19.8 (0.1)	2.2 (0.03)	0.3 (0.1)	1.7 (0.1)	0.2 (0.2)	0.2 (0.2)	0.1 (0.1)
A/S YA5 23	7.7 (0.0)	3.6 (0.04)	16.0 (1.0)	27.5 (0.2)	17.4 (0.3)	2.3 (0.1)	20.6 (0.3)	2.1 (0.01)	0.4 (0.1)	2.0 (0.4)	0.2 (0.2)	0.3 (0.1)	0.2 (0.01)
A/S YA5 31	7.9 (0.1)	3.6 (0.1)	15.7 (0.02)	27.7 (0.02)	17.5 (0.01)	2.3 (0.02)	20.3 (0.1)	2.2 (0.02)	0.4 (0.1)	1.8 (0.02)	0.2 (0.2)	0.3 (0.03)	0.2 (0.04)
A/S YA5 52	7.9 (0.02)	3.6 (0.01)	15.7 (0.1)	27.6 (0.04)	17.7 (0.1)	2.3 (0.04)	20.3 (0.2)	2.2 (0.02)	0.4 (0.1)	1.9 (0.2)	0.1 (0.1)	0.1 (0.1)	0.1 (0.1)
A/S YA5 95	8.0 (0.0)	3.7 (0.1)	15.5 (0.2)	27.6 (0.3)	17.7 (0.1)	2.3 (0.04)	20.3 (0.4)	2.2 (0.1)	0.5 (0.1)	1.9 (0.3)	0.2 (0.2)	0.2 (0.2)	0.1 (0.1)
A/S YA5 104	8.0 (0.01)	3.8 (0.0)	15.6 (0.02)	27.7 (0.4)	17.8 (0.02)	2.4 (0.0)	20.3 (0.04)	2.1 (0.0)	0.4 (0.1)	1.7 (0.03)	0.1 (0.1)	0.2 (0.2)	0.1 (0.1)

5 \* ± SD (n=2)

*Analysis of Flowering Times of A. thaliana Anti-Sense PDHK and pBI121 Control Transgenic Plants*

The anti-sense PDHK transgenic plants displayed a significantly earlier transition from the vegetative to the generative phase of growth, i.e. earlier initiation of the generative (flower formation) phase (recorded by monitoring the time, as days after planting: d.a.p.) compared to non-transformed wild-type and pBI121 plasmid only controls. As shown in Figure 11, 30-50% of the anti-sense PDHK transgenics were flowering as early as 31 d.a.p. compared to only 1-4% in the controls. This early flowering phenotype was even more dramatic at 34 d.a.p. when 50-75% of the anti-sense PDHK plants were in the generative phase compared to only 4-8% of the control plants. Most of the anti-sense PDHK plants were fully flowering (90% or greater floral initiation) by 39 d.a.p., but the non-transformed control plants and pBI121 plasmid only controls plants did not reach this stage until 46 d.a.p.

The time to reach maturity was also shorter in the anti-sense PDHK *A. thaliana* transgenics. For example, at 68 days after planting, all of the anti-sense PDHK transgenic lines had fully developed siliques and more than half of these were browned and mature. The few flowers that remained were senescing by this time. In contrast, the non-transformed wild type control plants and pBI121 control transformants still had significant flower development, primarily green immature siliques and only a few siliques which were brown and mature at this time. Under the growth conditions used by the inventors, the difference in maturity time was about 68-70 days for the anti-sense PDHK transgenic lines, compared to about 75-77 days for the control plants.

Given that the generation time in *Arabidopsis* control plants is about 75 days under the growth conditions used by the inventors, a 5 to 8-day earlier flowering and earlier maturing phenotype in the anti-sense PDHK



plants represents a shortening of generation time by about 10%. Similar modification of flowering time to extend the geographical range of cultivation is an important goal for *Brassica* crops (Lagercrantz et al., 1996). In related Brassicaceae (e.g. Canola), this would allow the advantage of an earlier harvest (e.g. on the Canadian Prairies) and permit more northerly cultivation (Murphy and Scarth, 1994). Late season frost damage in temperate climates could be avoided by earlier maturity, and this could also significantly alleviate problems associated with late-season clearing of chlorophyll from the maturing seeds (which can lead to "green oil" during processing and necessitate expensive bleaching steps).

The inventors' data demonstrates that increased respiration can hasten the transition from the vegetative to the generative phase of plant growth. It is interesting to note that the opposite effect, i.e. a *delay* in flowering time, was observed in transgenic plants in which citrate synthase was down-regulated by anti-sense technology, which resulted in a *decreased rate of respiration* in vegetative tissues (Landschütze et al., 1995). Thus, flowering time can be hastened by increased respiration and delayed by reduced respiration.

#### 20 *Analysis of Vegetative Growth of A.thaliana Anti-Sense PDHK and pBI121 Control Transgenic Plants*

The earlier-flowering phenotype of the anti-sense PDHK transgenics was correlated with an altered pattern of vegetative growth. There was a reduced accumulation of vegetative shoot tissue mass (Figure 12) correlated with a reduced number of rosette leaves produced in the anti-sense PDHK transgenics by the time plants switched to the generative (floral initiation) phase of growth (Figure 13).

To summarize, *A. thaliana* lines transformed with a constitutively-expressed anti-sense PDHK construct (designated as YA5 lines) exhibit both

altered vegetative growth and early flowering phenotypes, compared to the non-transformed controls or transformants containing only the selectable marker gene (transformed with pBI121), but without anti-sense PDHK. The difference with respect to the altered vegetative growth pattern phenotype of the anti-sense PDHK (YA5) transgenics (smaller plantlets with fewer rosette leaves compared to n-WT and pBI121 controls) was clearly visible at about 3.5 weeks after planting, and even more evident about one week later (30-31 days after planting). By about 31 days after planting, the early flowering phenotype was also apparent in the YA5 anti-sense PDHK transgenic lines.

Many of the plants were beginning to bolt or show visible floral meristem (flower bud) initiation, while there was no evidence of such development in the n-WT and pBI121 controls. At 40 to 42 days after planting, the early flowering phenotype of the YA5 anti-sense PDHK transgenic lines was very apparent, with most or all transgenics fully bolted with open flowers, while the n-WT and pBI121 controls show a much lower frequency of bolting.

Whereas flowering and generation time were shortened in the YA5 lines, the average number of siliques, seed weight and oil content were not adversely affected. Rather, as shown in Table 1 and Fig. 10, both average seed weight and fatty acid/oil content per seed were enhanced in the YA5 lines.

## EXPERIMENTAL PROCEDURES

### *General Molecular Biological Techniques:*

Isolation of plasmid DNA, restriction digestions, modification and ligation of DNA, PCR, agarose and polyacrylamide gel electrophoresis, transformation and culture of *E. coli* strains, DNA gel blot analyses (Southern, 1975) and RNA gel blot analyses, were carried out according to standard procedures as outlined by Sambrook et al., (1989).

### 10 *Cloning of YA5:*

An *Arabidopsis thaliana* (ecotype Columbia)  $\lambda$ YES cDNA expression library (Elledge et al., 1991) was obtained from Dr. Ronald Davis (Dept. of Biochemistry, Stanford University School of Medicine, Stanford CA 94305). Plasmids were generated by automatic subcloning procedures as described by Elledge et al. (1991). A putative *Escherichia coli* lyso-phosphatidic acid acyltransferase (LPAT, EC 2.3.1.51) mutant, JC201 (Coleman 1990) was obtained from Dr. Jack Coleman (Dept. of Biochemistry and Molecular Biology, Louisiana State University Medical Center, New Orleans, LA 70112). JC201 was transformed with the plasmids generated from the  $\lambda$ YES library, and selected at the non-permissive temperature of 44°C (Coleman, 1992). The YA5 cDNA from the temperature-insensitive transformant was sequenced on an Applied Biosystems Model 373A DNA Sequencing System using the Taq DyeDeoxy™ Terminator Cycle Sequencing Kit (Applied Biosystems, Inc.). The nucleotide and deduced amino acid sequences of clone YA5 were compared with sequences available in data banks using the FASTA program (Pearson and Lipman, 1988).

*Expression of the YA5 protein in E. coli:*

The YA5 full-length cDNA (YA5F) was cloned in pBluescript SK (+/-) in a 5' to 3' orientation of T7-T3. A primer encompassing the putative 5 translational initiation site Ompdk (AGGAGAGACTCGAGGCTTATGGCAGTGAAG) [SEQ ID NO:6] was synthesized to include an XhoI restriction site. Primer Ompdk and a T3 primer were used in a PCR reaction to amplify the YA5 encoding region from YA5F. The resulting PCR fragment was digested with HindIII (HindIII site is 10 present 3' to the stop codon) and XhoI, and cloned into pTrcHisB vector (Clontech) to generate construct pZTa5. SDS-PAGE analysis with lysates from IPTG-induced *E. coli* containing pZTa5 and pTrcHisB control vector confirmed that a novel protein of approximately 45 kDa was synthesized.

*15 Protein Kinase Assay with YA5-Expressing E. coli lysate:*

Protein kinase activity of the *E. coli* expressed YA5 protein was assayed essentially the same as described (Liu et al., 1995). The protein phosphorylation substrates, human  $E_1\alpha$  and  $E_1\beta$  PDC subunits, co-expressed and purified from *E. coli* M15, were obtained from Dr. Mulchand 20 S. Patel of the Department of Biochemistry, School of Medicine and Biomedical Sciences, State University of New York at Buffalo, Buffalo, New York. This co-expressed  $E_1\alpha$  and  $E_1\beta$  system has been used extensively in the study of the regulation of PDC  $E_1$  phosphorylation in mammalian systems (Korotchkina and Patel, 1995). For phosphorylation experiments, 20-25  $\mu$ g 25 of  $E_1\alpha/\beta$  was combined with about 10  $\mu$ g of YA5-accumulating *E. coli* cytosol protein, in a final volume of 100  $\mu$ l containing 20 mM potassium phosphate, pH 7.0, 1mM magnesium chloride, 2 mM dithiothreitol, 0. mM EDTA and 200  $\mu$ M cold ATP. The mixture was pre-incubated at room temperature for 5 minutes, and then 5  $\mu$ Ci  $^{32}$ P- $\gamma$ -ATP was added to start the assay. At 2, 5, 30 10, 15, and 20 minutes after starting the assay, 20  $\mu$ l aliquots of the reaction mixture were withdrawn and the reaction was stopped with 20  $\mu$ l of SDS-

denaturing mixture. Samples were then separated on 10% SDS-PAGE and autoradiographed to reveal  $^{32}\text{P}$ -labeled proteins.

*Construction of YA5 Antisense Plant Transformation Vector for Constitutive Expression:*

The YA5 cDNA contains internal BamHI (nt 628) and NcoI (nt1176) restriction sites. The BamHI and NcoI fragment was freed from YA5F and cloned into the respective sites in pBI524 (Datla et al., 1993), in an antisense orientation, under the control of a tandem 35S promoter. The YA5 antisense cassette was then excised from pBI524 by HindIII and EcoRI, and cloned into plant transformation vector pRD400 (Datla et al., 1992).

*Construction of YA5 Antisense and Partial Sense Plant Transformation Vectors for Seed-Specific Expression*

15 The YA5 full-length cDNA (YA5F; 1.5 kb ) was cloned in pBluescript SK (+/-) plasmid (Stratagene) in a 5' to 3' orientation of T7-T3. An 875 bp fragment was excised by a BamHI (Pharmacia) digestion and ligated into the plasmid pDH1 which had previously been cut with BamHI and dephosphorylated (treated with 1/10 unit of calf intestine alkaline  
20 phosphatase for 1 hour at 37°C). Plasmid pDH1 (provided by Dr. P. Covello, PBI/NRC) is the plasmid PE35SNT which has been manipulated so that the constitutive tandem 35S promoter has been excised and replaced with the seed-specific napin promoter, obtained from plasmid pUC19. Ligations were performed at 4-12 °C overnight in a waterbath, following the instructions  
25 provided by the manufacturer. Competent *E. coli* cells (DH5 $\alpha$ , Gibco BRL) were transformed by a heat shock method, with 50-100 ng of transforming DNA, plated on a selective medium (LB with 50  $\mu\text{g}/\text{mL}$  ampicillin) and incubated overnight at 37 °C. The Bluescript plasmid DNA (10 ng) was used as a positive control for the transformation. Single transformed cells were  
30 grown overnight (37 °C, 225 r.p.m.) in 5 mL LB with 50  $\mu\text{g}/\text{mL}$  ampicillin.

DNA extraction and purification was performed with a Qiaprep Spin Miniprep kit (Qiagen). Restriction digestions were performed with HindIII to check for the presence and the orientation of inserts in the plasmid. In the case of a YA5 insert in an anti-sense orientation, two fragments of about 1.0 and 1.4 5 Kb were obtained, while a YA5 insert in a sense orientation gave two fragments of about 1.8 and 0.6 Kb. The cassette and the insert (in either sense or anti-sense orientation) were excised by a partial double digestion with HindIII and EcoRI (1unit/20  $\mu$ L reaction for 10 min at 37 °C). The DNA fragments corresponding to the cassette with either the sense or anti-sense 10 insert, were purified from the agarose gel using a Geneclean II Kit (Bio 101, Inc.) and ligated to the HindIII/EcoRI digested pRD400 plasmid. The best ligation results were obtained with a 1:10 plasmid to insert ratio in a 10  $\mu$ L reaction volume, using 1  $\mu$ L of T4 ligase and buffer (New England Biolabs) at 4 °C overnight. The reaction mixture was heated at 45 °C for 5 min and ice 15 chilled prior to adding the ligase. The following day, 1  $\mu$ L of T4 ligase was added and the mixture left at room temperature for a further 3-4 hrs. The identity of each construct was re-checked by digestion and DNA sequencing before plant transformation.

#### 20 *Transformation of Arabidopsis thaliana:*

The transformation protocol was adapted from that described by Bechtold et al., (1993). Plants of *Arabidopsis thaliana* ecotype Columbia were grown in moist soil at a density of 10-12 plants per pot, in 4-inch square pots, and covered with a nylon screen fixed in place with an elastic band. 25 Once the plants reached the stage at which bolts were just emerging, plants were watered, the bolts and some of the leaves were clipped, and the plants infiltrated in *Agrobacterium* suspension as outlined below.

To grow the *Agrobacterium*, a 25 mL suspension in LB medium containing kanamycin at a concentration of 50  $\mu$ g/mL was cultured for two to 30 three days ahead of time. The day before infiltration, this “seed culture” was

added to 400 mL of LB medium containing 50  $\mu\text{g/mL}$  kanamycin. Once the absorbance at 600 nm was  $> 2.0$ , the cells were harvested by centrifugation (5,000  $\times g$ , 10 min in a GSA rotor at room temperature) and resuspended in 3 volumes of infiltration medium (1/2 x Murashige and Skoog salts, 1 x B5 5 vitamins, 5.0% sucrose, 0.044  $\mu\text{M}$  benzylaminopurine) to an optical density at 600 nm of 0.8. The *Agrobacterium* suspension was then poured into a beaker and the potted plants inverted into the beaker so that the bolts and entire rosettes were submerged. The beaker was then placed into a large Bell jar and a vacuum drawn using a vacuum pump, until bubbles formed on 10 the leaf and stem surfaces and the solution started to bubble a bit, and then the vacuum was released rapidly. [Note: The necessary time and pressure will vary from one lab setup to the next, but good infiltration is visibly apparent as uniformly darkened, water-soaked tissue.] Pots were removed from the beaker, laid on their side in a plastic tray and covered with a plastic 15 dome, to maintain humidity. The following day, the plants were uncovered, set upright and allowed to grow for approximately four weeks in a growth chamber under continuous light conditions as described by Katavic et al., (1995). When the siliques were mature and dry, seeds were harvested and selected for positive transformants.

20

*Selection of Putative Transformants (Transgenic plants) and Growth and Analysis of Transgenic Plants:*

Seeds harvested from vacuum-infiltration transformation experiments were sterilized by treating for 1 min in ethanol and then 5 min in 25 50% bleach/0.05% Tween 20™ in sterile distilled water. Then the seeds were rinsed several times with sterile distilled water. Seeds were plated by resuspending them in sterile 0.1% agarose at room temperature (about 1 mL agarose for every 500-1000 seeds), and then applying a volume equivalent to about 2,000-4,000 seeds onto 150 x 15 mm selection plates (1/2 x 30 Murashige and Skoog salts, 0.8 % agar, autoclave, cool and add 1 x B5

vitamins and kanamycin at a final concentration of 50  $\mu\text{g}/\text{mL}$ ). The plates were dried in a laminar flow hood until seed no longer flowed when the plates were tipped. The plates were then vernalized for two nights at 4°C in the dark, and then moved to a growth chamber (conditions as described by 5 Katavic et al., 1995). After 7-10 days, transformants were clearly identifiable as dark green plants with healthy green secondary leaves and roots that extended over and into the selective medium.

Seedlings were transplanted to soil, plants grown to maturity and mature seeds ( $T_2$  generation as defined in Katavic et al., 1994) collected and 10 analyzed.  $T_2$  seeds were propagated. The vegetative growth patterns were monitored by measuring shoot tissue dry weights, and/or by counting the number of rosette leaves present by the time plants began to enter the generative (flower initiation) stage. Floral initiation (beginning of generative phase of growth) was analyzed by recording, on a daily basis, the 15 percentage of plants in which a flower bud first appeared and/or the percentage of plants that were bolting (as described by Zhang et al. 1997). Data was reported in terms of percentage of plants flowering/bolting on a given day after planting (d.a.p.).

#### 20 *Preparation of Mitochondria from A. thaliana Non-Transformed Control and Anti-YA5 Transgenic Plants*

All extractions were performed at 4°C. An enriched mitochondrial fraction was prepared by a procedure modified from Ap Rees et al., (1993). About 30 g of freshly-harvested shoots were homogenized using a chilled 25 Waring Bleder in 4 volumes of extraction buffer (50 mM Tris-HCl, pH 8.0, containing 300 mM mannitol, 5mM EDTA, 0.1 % bovine serum albumin, 1% PVPP (polyvinylpolypyrrolidone) and 9 mM 2-mercaptoethanol. The homogenate was filtered through four layers of cheesecloth and one layer of Miracloth and centrifuged at 2,000  $\times g$  for 10 min. The pellet was discarded 30 and the supernatant centrifuged at 10,000  $\times g$  for 30 min. The pellet was



resuspended in the extraction buffer minus PVPP and used as an "enriched" mitochondrial preparation. For the succinate dehydrogenase activity measurements, this preparation was used directly. For measurements of PDC, fumarase and citrate synthase, the freshly-prepared mitochondria were first lysed in the presence of 0.1% (v/v) Triton X-100 to release the mitochondrial enzymes. The Triton lysate was clarified by centrifugation at 27,000 x g and the supernatant, containing solubilized enzymes, was concentrated with a Centricon-30 filter concentrator (Amicon). The resulting concentrate was used as the enzyme source in a PDC assay. Protein concentrations were estimated by the method of Bradford (1976) using bovine serum albumin as a standard, and normalized prior to assay.

*Pyruvate Dehydrogenase Complex Assay:*

The method used to determine the pyruvate dehydrogenase (PDH) complex activity present in mitochondrial protein preparations was modified from the method of Reid et al., (1977). The assay mixture consisted of 0.1 mM TPP, 5 mM MgCl<sub>2</sub>, 1.5 mM NAD<sup>+</sup>, 0.1 mM Coenzyme A, 3.0 mM cysteine-HC and 1.5 mM pyruvate in 100 mM Tricine pH 8.0, in a final volume of 2 mL. Reactions were initiated by the addition of mitochondrial lysate concentrate. Control reactions contained all components except pyruvate. The reaction mixture was incubated at 30°C and the formation of NADH was monitored at a wavelength of 340 nm at 15-second intervals for 3 minutes, using a Beckman DU 74 spectrophotometer.

*25 Citrate Synthase Assay:*

The method used to measure citrate synthase activity present in mitochondrial protein preparations was modified from the method of Srere (1969). The reaction mixture contained 0.2 mM 5',5'-dithiobis-2-nitrobenzoic acid (DTNB), 0.1 mM acetyl-CoA, and mitochondrial lysate protein, in 50 mM Tris-HCl, pH 7.8, in a final reaction volume of 2 mL, incubated at 30°C. The

absorption at 412 nm was followed for 3 minutes to measure possible acetyl-CoA deacylase activity. The citrate synthase reaction was then started by the addition of 0.5 mM oxaloacetate (OAA), and the release of Coenzyme A (CoASH), the SH group of which reacts with the DTNB (Ellman's reagent), was monitored at 412 nm. The resulting mercaptide ion has a strong absorption ( $E = 13,600$ ) at 412 nm. Control reactions contained all components except OAA.

*Fumarase Assay:*

10 Fumarase activity present in mitochondrial protein preparations was assayed by the method of Hill and Bradshaw (1969). The reaction mixture contained 25 mM sodium malate and mitochondrial lysate protein in 50 mM sodium phosphate buffer, pH 7.5, in a final reaction volume of 1 mL. Reactions were incubated at 28°C. Fumarase activity, measured by the  
15 formation of fumarate, was determined spectrophotometrically by monitoring the increase in absorbance at 250 nm, at 15-second intervals for two minutes.

*Succinate Dehydrogenase Assay:*

20 Mitochondrial succinate dehydrogenase activity was measured using the method of Veeger et al., (1969). The reaction mixture contained 1 mM KCN, 40 mM succinate, 1 mM EDTA, 0.1% BSA, 3 mM  $K_3Fe(CN)_6$ , 0.1% Triton X-100 and unlysed mitochondria in 100 mM sodium phosphate buffer at pH 7.5, in a final reaction volume of 1 mL. Reaction mixtures were  
25 incubated at 28°C. The reaction was initiated by the addition of mitochondria in oxygen-free phosphate buffer. The change in absorbance at 455 nm was monitored spectrophotometrically, at 20-second intervals for 2 minutes. Control reactions contained all components except succinate.

*Seed Lipid Analyses of Seeds from A. thaliana Non-Transformed Controls and Anti-Sense PDHK (AsYA5) Transgenics:*

Mature siliques and T<sub>2</sub> or T<sub>3</sub> seeds were isolated from anti-sense PDHK transformants (designated as YA5 lines) or pBI121 control 5 transformants (without anti-sense PDHK, but with a kanamycin resistance gene) and siliques and seeds were also isolated from non-transformed wild-type control plants and their respective seed oil contents, fatty acyl compositions of the seed oils, average seed weights, number of siliques per 15 cm segment of bolting stem and number of seeds per silique were 10 determined as described by Zou et al., (1997). All plants were grown in the same growth chamber, at the same time and under identical light and temperature regimes, as described previously (Katavic et al., 1995; Zou et al., 1997). Because of the extremely small seed size and weight, analyses were performed on 100- or 400-seed replicates that were carefully counted 15 out under a dissecting microscope. Seed samples were ground using a polytron in chloroform:isopropanol (2:1, [v/v]) containing 0.2% w/v butylated hydroxytoluene and tripentadecanoin as an internal standard. All other conditions for the isolation and analysis of seed total fatty acid content and fatty acid composition (expressed as wt% of total fatty acids) by gas 20 chromatography were performed as detailed previously (Taylor et al., 1992b; Taylor et al., 1995; Katavic et al., 1995, Zou et al., 1997). Oil content was expressed either as µg of total fatty acids per sample size (See Fig. 10), or as mg of oil per sample size (See Table 1), calculated by assuming 3 moles of fatty acids per mole of triacylglycerol (oil), as described by Zou et al., 25 (1997).

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

(A) NAME: National Research Council of Canada  
(B) STREET: 1200 Montreal Road  
(C) CITY: Ottawa  
(D) STATE: Ontario  
(E) COUNTRY: Canada  
(F) POSTAL CODE (ZIP): K1A 0R6

(ii) TITLE OF INVENTION: Plant Pyruvate Dehydrogenase Kinase Gene

(iii) NUMBER OF SEQUENCES: 9

## (iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

## (v) CURRENT APPLICATION DATA:

APPLICATION NUMBER: WO PCT/CA98/-----

## (vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 60/038,815  
(B) FILING DATE: 10-FEB-1997

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1457 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Arabidopsis thaliana

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

TCCATCTGCG CACTTCTTTC GTCCAGTCCA TGATAATAAC GGTGGAGAAC GACGGAGGCG	60
GGCGACGTTA GGGTTTCTAA TCATTCTCT CTCTTAGACG CTTATGGCAG TGAAGAAAGC	120
CTGCGAAATG TTCCCGAAGA GTTTGATCGA AGATGTTTAC AAATGGGGTT GCATGAAGCA	180
AACCGGTGTT AGCCTTAGAT ACATGATGGA GTTTGGTTCC AAACCTACTG AGAGGAATCT	240
TTTGATTCTT GCTCAGTTTT TGCATAAGGA GCTTCCGATT CGCGTCGCCA GGAGAGCGAT	300

```

CGAACTCCAG ACGCTTCCTT ATGGTCTCTC TGATAAACCT GCCGTTTGA AGGTGCGGGA      360
TTGGTATTTG GAATCTTTCA GGGACATGAG AGCATTTCCT GAGATTAAGG ATTCGGGTGA      420
CGAGAAAGAT TTCACTCAGA TGATTAAGGC TGTCAAAGTA AGGCATAACA ATGTGGTTCC      480
CATGATGGCT TTGGGTGTTA ATCAGCTCAA GAAAGGAATG AATTCTGGAA ATCTTGATGA      540
GATTCATCAG TTTCTTGATC GTTCTACTT GTCGCGAATC GGGATCCGGA TGCTTATTGG      600
GCAGCACGTT GAGTTGCATA ATCCAAATCC ACCGCTTCAT ACAGTGGGTT ATATACACAC      660
AAAGATGTCT CCTATGGAGG TAGCAAGGAA TGCAAGTGAA GATGCTCGGT CAATTTGTTT      720
CCGAGAGTAC GGTTCCTGCAC CGGAAATAAA CATATATGGC GATCCCAGTT TCACCTTTCC      780
GTATGTTCCA ACGCATTTGG ATCTTATGAT GTATGAGCTA GTCAAGAACT CTCTACGTGC      840
TGTCCAAGAG CGATTTGTTG ACTCTGATAG AGTTGCACCA CCAATCCGCA TTATAGTTGC      900
TGATGGAATC GAAGATGTTA CTATAAAGGT CTCAGATGAA GGTGGAGGTA TAGCAAGAAG      960
CGGTCTTCCC AGAATATTCA CCTATCTTTA CAGCACTGCA AGAAACCCGC TTGAGGAGGA     1020
TGTCGATTTA GGAATAGCTG ATGTTCCCGG GACTATGGGT GGATATGGTT ATGGTCTTCC     1080
AATTAGTCGC TTGTATGCTC GATATTTCCG TGGAGATTTG CAGATCATAT CCATGGAAGG     1140
ATATGGGACT GATGCATACT TGCACTTGTC TCGCCTTGA GATTCGCAAG AGCCTTTACC     1200
CTGAGAACAT CTCTATGTCA GGCAAAGTAA AGAAAGCTTT GACATGTATT TATGGTAGAT     1260
GAGGGATATC TACAATACTC AATTATTTAT GCTTTTCCAG TTTCTGCTAA TGTACAGACT     1320
ACAGACATTA TTTTCTCGTA TTACGCTTTC TTGATTTTAG ACTCAGATAT GGAGCTTTTT     1380
CCAAGTGAGT TAATCTCCTA TGATTGTTT TGGTTCGATC CAAAACCACC TTGTATCCGA     1440
AAAAAAAAAA AAAAAAA                                     1457

```

## (2) INFORMATION FOR SEQ ID NO: 2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 366 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (iii) HYPOTHETICAL: YES

## (iv) ANTI-SENSE: NO

## (v) FRAGMENT TYPE: N-terminal

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Arabidopsis thaliana*

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met	Ala	Val	Lys	Lys	Ala	Cys	Glu	Met	Phe	Pro	Lys	Ser	Leu	Ile	Glu	1	5	10	15
Asp	Val	His	Lys	Trp	Gly	Cys	Met	Lys	Gln	Thr	Gly	Val	Ser	Leu	Arg	20	25	30	
Tyr	Met	Met	Glu	Phe	Gly	Ser	Lys	Pro	Thr	Glu	Arg	Asn	Leu	Leu	Ile	35	40	45	
Ser	Ala	Gln	Phe	Leu	His	Lys	Glu	Leu	Pro	Ile	Arg	Val	Ala	Arg	Arg	50	55	60	
Ala	Ile	Glu	Leu	Gln	Thr	Leu	Pro	Tyr	Gly	Leu	Ser	Asp	Lys	Pro	Ala	65	70	75	80
Val	Leu	Lys	Val	Arg	Asp	Trp	Tyr	Leu	Glu	Ser	Phe	Arg	Asp	Met	Arg	85	90	95	
Ala	Phe	Pro	Glu	Ile	Lys	Asp	Ser	Gly	Asp	Glu	Lys	Asp	Phe	Thr	Gln	100	105	110	
Met	Ile	Lys	Ala	Val	Lys	Val	Arg	His	Asn	Asn	Val	Val	Pro	Met	Met	115	120	125	
Ala	Leu	Gly	Val	Asn	Gln	Leu	Lys	Lys	Gly	Met	Asn	Ser	Gly	Asn	Leu	130	135	140	
Asp	Glu	Ile	His	Gln	Phe	Leu	Asp	Arg	Phe	Tyr	Leu	Ser	Arg	Ile	Gly	145	150	155	160
Ile	Arg	Met	Leu	Ile	Gly	Gln	His	Val	Glu	Leu	His	Asn	Pro	Asn	Pro	165	170	175	
Pro	Leu	His	Thr	Val	Gly	Tyr	Ile	His	Thr	Lys	Met	Ser	Pro	Met	Glu	180	185	190	
Val	Ala	Arg	Asn	Ala	Ser	Glu	Asp	Ala	Arg	Ser	Ile	Cys	Phe	Arg	Glu	195	200	205	
Tyr	Gly	Ser	Ala	Pro	Glu	Ile	Asn	Ile	Tyr	Gly	Asp	Pro	Ser	Phe	Thr	210	215	220	
Phe	Pro	Tyr	Val	Pro	Thr	His	Leu	Asp	Leu	Met	Met	Tyr	Glu	Leu	Val	225	230	235	240
Lys	Asn	Ser	Leu	Arg	Ala	Val	Gln	Glu	Arg	Phe	Val	Asp	Ser	Asp	Arg	245	250	255	
Val	Ala	Pro	Pro	Ile	Arg	Ile	Ile	Val	Ala	Asp	Gly	Ile	Glu	Asp	Val	260	265	270	
Thr	Ile	Lys	Val	Ser	Asp	Glu	Gly	Gly	Gly	Ile	Ala	Arg	Ser	Gly	Leu	275	280	285	
Pro	Arg	Ile	Phe	Thr	Tyr	Leu	Tyr	Ser	Thr	Ala	Arg	Asn	Pro	Leu	Glu	290	295	300	

45

Glu Asp Val Asp Leu Gly Ile Ala Asp Val Pro Gly Thr Met Gly Gly  
 305 310 315 320

Tyr Gly Tyr Gly Leu Pro Ile Ser Arg Leu Tyr Ala Arg Tyr Phe Gly  
 325 330 335

Gly Asp Leu Gln Ile Ile Ser Met Glu Gly Tyr Gly Thr Asp Ala Tyr  
 340 345 350

Leu His Leu Ser Arg Leu Gly Asp Ser Gln Glu Pro Leu Pro  
 355 360 365

## (2) INFORMATION FOR SEQ ID NO: 3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 405 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: N-terminal

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Porcine PDH kinase subunit I

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Met Arg Trp Phe Arg Ala Leu Leu Lys Asn Ala Ser Leu Ala Gly Ala  
 1 5 10 15

Pro Lys Tyr Ile Glu His Phe Ser Lys Phe Ser Pro Ser Pro Leu Ser  
 20 25 30

Met Lys Gln Phe Leu Asp Phe Gly Ser Ser Asn Ala Cys Glu Lys Thr  
 35 40 45

Ser Phe Thr Phe Leu Arg Gln Glu Leu Pro Val Arg Leu Ala Asn Ile  
 50 55 60

Met Lys Glu Ile Asn Leu Leu Pro Asp Arg Val Leu Ser Thr Pro Ser  
 65 70 75 80

Val Gln Leu Val Gln Ser Trp Tyr Val Gln Ser Leu Leu Asp Ile Met  
 85 90 95

Glu Phe Leu Asp Lys Asp Pro Glu Asp His Arg Thr Leu Ser Gln Phe  
 100 105 110

Thr Asp Ala Leu Val Thr Ile Arg Asn Arg His Asn Asn Val Val Pro  
 115 120 125

## 46

```

Thr Met Ala Gln Gly Val Leu Glu Tyr Lys Asp Thr Tyr Gly Asp Asp
130                               135                               140

Pro Val Ser Asn Gln Asn Ile Gln Tyr Phe Leu Asp Arg Phe Tyr Leu
145                               150                               155                               160

Ser Arg Ile Ser Ile Arg Met Leu Ile Asn Gln Thr Leu Ile Phe Asp
165                               170                               175

Gly Ser Thr Asn Pro Ala His Pro Lys His Gly Ser Ile Asp Pro Asn
180                               185                               190

Pro Asn Cys Ser Val Ser Asp Val Val Lys Asp Ala Tyr Asp Met Ala
195                               200                               205

Lys Leu Leu Cys Asp Lys Tyr Tyr Met Ala Ser Pro Asp Leu Glu Ile
210                               215                               220

Gln Glu Val Asn Ala Thr Asn Ala Thr Gln Pro Ile His Met Val Tyr
225                               230                               235                               240

Val Pro Ser His Leu Tyr His Met Leu Phe Glu Leu Phe Lys Asn Ala
245                               250                               255

Met Arg Ala Thr Val Glu Ser His Glu Ser Ser Leu Thr Leu Pro Pro
260                               265                               270

Ile Lys Ile Met Val Ala Leu Gly Glu Glu Asp Leu Ser Ile Lys Met
275                               280                               285

Ser Asp Arg Gly Gly Gly Val Pro Leu Arg Lys Ile Glu Arg Leu Phe
290                               295                               300

Ser Tyr Met Tyr Ser Thr Ala Pro Thr Pro Gln Pro Gly Thr Gly Gly
305                               310                               315                               320

Thr Pro Leu Ala Gly Phe Gly Tyr Gly Leu Pro Ile Ser Arg Leu Tyr
325                               330                               335

Ala Lys Tyr Phe Gln Gly Asp Leu Gln Leu Phe Ser Met Glu Gly Phe
340                               345                               350

Gly Thr Asp Ala Val Ile Tyr Leu Lys Ala Leu Ser Thr Asp Ser Val
355                               360                               365

Glu Arg Leu Pro Val Tyr Asn Lys Ser Ala Trp Arg His Tyr Gln Thr
370                               375                               380

Ile Gln Glu Ala Gly Asp Trp Cys Val Pro Ser Thr Glu Pro Lys Asn
385                               390                               395                               400

Thr Tyr Arg Val Ser
405

```

## (2) INFORMATION FOR SEQ ID NO: 4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 434 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single



47

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: N-terminal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Porcine PDH kinase subunit II

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

```

Met Arg Leu Ala Arg Leu Leu Arg Gly Gly Thr Ser Val Arg Pro Leu
 1           5           10           15

Cys Ala Val Pro Cys Ala Ser Arg Ser Leu Ala Ser Asp Ser Ala Ser
      20           25           30

Gly Ser Gly Pro Ala Ser Glu Ser Gly Val Pro Gly Gln Val Asp Phe
 35           40           45

Tyr Ala Arg Phe Ser Pro Ser Pro Leu Ser Met Lys Gln Phe Leu Asp
 50           55           60

Phe Gly Ser Val Asn Ala Cys Glu Lys Thr Ser Phe Met Phe Leu Arg
 65           70           75           80

Gln Glu Leu Pro Val Arg Leu Ala Asn Ile Met Lys Glu Ile Ser Leu
      85           90           95

Leu Pro Asp Asn Leu Leu Arg Thr Pro Ser Val Gln Leu Val Gln Ser
      100           105           110

Trp Tyr Ile Gln Ser Leu Gln Glu Leu Leu Asp Phe Lys Asp Lys Ser
      115           120           125

Ala Glu Asp Ala Lys Thr Ile Tyr Glu Phe Thr Asp Thr Val Ile Arg
      130           135           140

Ile Arg Asn Arg His Asn Asp Val Ile Pro Thr Met Ala Gln Gly Val
      145           150           155           160

Asn Glu Tyr Lys Glu Ser Phe Gly Ser Asp Pro Val Thr Ser Gln Asn
      165           170           175

Val Gln Tyr Phe Leu Asp Arg Phe Tyr Met Ser Arg Ile Ser Ile Arg
      180           185           190

Met Leu Leu Asn Gln His Ser Leu Leu Phe Gly Gly Lys Gly Ser Pro
      195           200           205

Ser His Arg Lys His Ile Gly Ser Ile Asn Pro Asn Cys Asp Val Val
      210           215           220

Glu Val Ile Lys Asp Gly Tyr Glu Asn Ala Arg Arg Leu Cys Asp Leu
      225           230           235           240

```

[illegible]

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 412 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

## 49

Met Ile Leu Thr Ser Val Leu Gly Ser Gly Pro Arg Ser Gly Ser Ser  
 1 5 10 15  
 Leu Trp Pro Leu Leu Gly Ser Ser Leu Ser Leu Arg Val Arg Ser Thr  
 20 25 30  
 Ser Ala Thr Asp Thr His His Val Glu Leu Ala Arg Glu Arg Ser Lys  
 35 40 45  
 Thr Val Thr Ser Phe Tyr Asn Gln Ser Ala Ile Asp Val Val Ala Glu  
 50 55 60  
 Lys Pro Ser Val Arg Leu Thr Pro Thr Met Met Leu Tyr Ser Gly Arg  
 65 70 75 80  
 Ser Gln Asp Gly Ser His Leu Leu Lys Ser Gly Arg Tyr Leu Gln Gln  
 85 90 95  
 Glu Leu Pro Val Arg Ile Ala His Arg Ile Lys Gly Phe Val Val Phe  
 100 105 110  
 Leu Ser Ser Leu Val Ala Thr Leu Pro Tyr Cys Thr Val His Glu Leu  
 115 120 125  
 Tyr Ile Arg Ala Phe Gln Lys Leu Thr Asp Phe Pro Pro Ile Lys Asp  
 130 135 140  
 Gln Ala Asp Glu Ala Gln Tyr Cys Gln Leu Val Arg Gln Leu Leu Asp  
 145 150 155 160  
 Asp His Lys Asp Val Val Thr Leu Leu Ala Glu Gly Leu Arg Glu Ser  
 165 170 175  
 Arg Lys His Ile Glu Asp Glu Lys Leu Val Arg Tyr Phe Leu Asp Lys  
 180 185 190  
 Thr Leu Thr Ser Arg Leu Gly Ile Arg Met Leu Ala Thr His His Leu  
 195 200 205  
 Ala Leu His Glu Asp Lys Pro Asp Phe Val Gly Ile Ile Ser Thr Arg  
 210 215 220  
 Leu Ser Pro Lys Lys Ile Ile Glu Lys Trp Val Asp Phe Ala Arg Arg  
 225 230 235 240  
 Leu Cys Glu His Lys Tyr Gly Asn Ala Pro Arg Val Arg Ile Asn Gly  
 245 250 255  
 His Val Ala Ala Arg Phe Pro Phe Ile Pro Met Pro Leu Asp Tyr Ile  
 260 265 270  
 Leu Pro Glu Leu Leu Lys Asn Ala Met Arg Ala Thr Met Glu Ser His  
 275 280 285  
 Leu Asp Thr Pro Tyr Asn Val Pro Asp Val Val Ile Thr Ile Ala Asn  
 290 295 300  
 Asn Asp Val Asp Leu Ile Ile Arg Ile Ser Asp Arg Gly Gly Gly Ile  
 305 310 315 320

50

Ala His Lys Asp Leu Asp Arg Val Met Asp Tyr His Phe Thr Thr Ala  
325 330 335

Glu Ala Ser Thr Gln Asp Pro Arg Ile Ser Pro Leu Phe Asp His Leu  
340 345 350

Asp Thr His Ser Gly Gly Gln Ser Gly Pro Met His Gly Phe Gly Phe  
355 360 365

Gly Leu Pro Thr Ser Arg Ala Tyr Ala Glu Tyr Leu Gly Gly Ser Leu  
370 375 380

Gln Leu Gln Ser Leu Gln Gly Ile Gly Thr Asp Val Leu His Arg Ser  
385 390 395 400

Arg His Ile Asp Gly Arg Glu Glu Ser Phe Arg Ile  
405 410

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

AGGAGAGACT CGAGGCTTAT GGCAGTGAAG

30

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Arabidopsis thaliana*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

Gly Xaa Gly Xaa Gly  
1 5

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 5 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Arabidopsis thaliana

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Asp Xaa Gly Xaa Gly  
1 5

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 9 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Arabidopsis thaliana

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Glu Leu Xaa Lys Asn Xaa Xaa Arg Ala  
1 5



# American Type Culture Collection

12301 Parklawn Drive • Rockville, MD 20852 USA • Telephone: 301-231-5519 or 231-5532 • FAX: 301-816-4344

BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF  
THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT PROCEDURE

## INTERNATIONAL FORM

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT ISSUED PURSUANT TO RULE 7.3  
AND VIABILITY STATEMENT ISSUED PURSUANT TO RULE 10.2

To: (Name and Address of Depositor or Attorney)

National Research Council of Canada  
Attn: Dr. David C. Taylor  
Plant Biotechnology Institute, 110 Gymnasium Place  
Saskatoon, SK S7N 0W9  
Canada

Deposited on Behalf of: National Research Council of Canada

Identification Reference by Depositor: ATCC Designation

pAsYA5 plasmid 209561

The deposits were accompanied by: \_\_\_ a scientific description \_\_\_ a proposed taxonomic description indicated above.

The deposits were received December 18, 1997 by this International Depository Authority and have been accepted.

### AT YOUR REQUEST:

☒ We will inform you of requests for the strains for 30 years.

The strains will be made available if a patent office signatory to the Budapest Treaty certifies one's right to receive, or if a U.S. Patent is issued citing the strains, and ATCC is instructed by the United States Patent & Trademark Office or the depositor to release said strains.

If the cultures should die or be destroyed during the effective term of the deposit, it shall be your responsibility to replace them with living cultures of the same.

The strains will be maintained for a period of at least 30 years from date of deposit, or five years after the most recent request for a sample, whichever is longer. The United States and many other countries are signatory to the Budapest Treaty.

The viability of the cultures cited above was tested December 30, 1997. On that date, the cultures were viable.

International Depository Authority: American Type Culture Collection, Rockville, Md. 20852 USA

Signature of person having authority to represent ATCC:

Barbara M. Hailey, Administrator, Patent Depository

Date: January 20, 1998

cc: Kirby, Eades, Gale and Baker



# American Type Culture Collection

12301 Parklawn Drive • Rockville, MD 20852 USA • Telephone: 301-231-5519 or 131-5532 • FAX: 301-816-4344

BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF  
THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT PROCEDURE

## INTERNATIONAL FORM

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT ISSUED PURSUANT TO RULE 7.3  
AND VIABILITY STATEMENT ISSUED PURSUANT TO RULE 10.2

To: (Name and Address of Depositor or Attorney)

National Research Council of Canada  
Attn: Dr. David C. Taylor  
Plant Biotechnology Institute, 110 Gymnasium Place  
Saskatoon, SK S7N 0W9  
Canada

Deposited on Behalf of: National Research Council of Canada

Identification Reference by Depositor: ATCC Designation

pYA5 plasmid 209562

The deposits were accompanied by: \_\_\_\_ a scientific description \_\_\_\_ a proposed taxonomic description indicated above.

The deposits were received December 18, 1997 by this International Depository Authority and have been accepted.

### AT YOUR REQUEST:

☒ We will inform you of requests for the strains for 30 years.

The strains will be made available if a patent office signatory to the Budapest Treaty certifies one's right to receive, or if a U.S. Patent is issued citing the strains, and ATCC is instructed by the United States Patent & Trademark Office or the depositor to release said strains.

If the cultures should die or be destroyed during the effective term of the deposit, it shall be your responsibility to replace them with living cultures of the same.

The strains will be maintained for a period of at least 30 years from date of deposit, or five years after the most recent request for a sample, whichever is longer. The United States and many other countries are signatory to the Budapest Treaty.

The viability of the cultures cited above was tested January 2, 1998. On that date, the cultures were viable.

International Depository Authority: American Type Culture Collection, Rockville, Md. 20852 USA

Signature of person having authority to represent ATCC:

Barbara M. Hailey, Administrator, Patent Depository

Date: January 20, 1998

cc: Kirby, Eades, Gale and Baker

REFERENCES RELEVANT TO THE CURRENT INVENTION:

- Ap Rees, T., Bryce, J.H., Wilson, P.M. and Green, J.H. (1983) Role and location of NAD malic enzyme in thermogenic tissue of *Araceae*. *Arch. Biochem. Biophys.* **227**: 511-521.
- Bechtold, N., Ellis, J. and Pelletier, G. (1993) *In planta Agrobacterium*-mediated gene transfer by infiltration of adult *Arabidopsis thaliana* plants. *C.R. Acad. Sci. Ser. III Sci. Vie*, **316**: 1194-1199.
- Becker, D., Brettschneider, R. and Lörz, H. (1994) Fertile transgenic wheat from microprojectile bombardment of scutellar tissue. *Plant J.* **5**: 299-307.
- Bradford, M.M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**: 248-254.
- Budde R. J. A., Fang, T.K. and Randall, D.D. (1988) Regulation of the phosphorylation of mitochondrial pyruvate dehydrogenase complex *in situ*. *Plant Physiol.*, **88**: 1031-1036.
- Coleman, J. (1990) Characterization of *Escherichia coli* cells deficient in 1-acyl-*sn*-glycerol-3-phosphate acyltransferase activity. *J. Biol. Chem.* **256**: 17215-17221.
- Coleman, J. (1992) Characterization of the *Escherichia coli* gene for 1-acyl-*sn*-glycerol-3-phosphate acyltransferase (*plsC*). *Mol. Gen. Genet.* **232**: 295-303.
- Damuni, Z., Merryfield, M.L., Humphreys, J.S. and Reed, L.J. (1984) Purification and properties of branched-chain  $\alpha$ -ketoacid dehydrogenase phosphatase from bovine kidney. *Proc. Natl. Acad. Sci. USA.* **81**: 4335-4338.
- Datta, R.S.S., Hammerlindl, J.K., Panchuk, B., Pelcher, L.E. and Keller, W.A. (1992) Modified binary plant transformation vectors with the wild-type gene encoding NPTII. *Gene* **211**: 383-384.



- Datla, R.S.S., Bekkaoui, F., Hammerlindl, J., Pilate, G., Dunstan, D.I. and Crosby, W.L. (1993) Improved high-level constitutive foreign gene expression in plants using an AMV RNA4 untranslated leader sequence. *Plant Sci.* **94**: 139-149.
- Datla, R., Anderson, J.W. and Selvaraj, G. (1997) Plant promoters for transgene expression. *Biotechnology Annual Review* **3**: 269-296.
- DeBlock, M., DeBrouwer, D. and Tenning P. (1989) Transformation of *Brassica napus* and *Brassica oleracea* using *Agrobacterium tumefaciens* and the expression of the *bar* and *neo* genes in the transgenic plants. *Plant Physiol.* **91**: 694-701.
- De Lange, P., Van Blokland, R., Kooter, J.M. and Mol, J.M.N. (1995) Suppression of flavenoid flower pigmentation genes in *Petunia hybrida* by the introduction of antisense and sense genes. *In: Gene Silencing in Higher Plants and Related Phenomena in Other Eukaryotes*. P.Meyer (Ed.), Springer-Verlag, Berlin, pp. 55-75.
- Elledge, S.J., Mulligan, J.T., Ramer, S.W., Spottswood, M. and Davis, R.W. (1991)  $\lambda$ YES: A multifunctional cDNA expression vector for the isolation of genes by complementation of yeast and *Escherichia coli* mutations. *Proc. Natl. Acad. Sci. USA*, **88**: 1731-1735.
- Gale, J. (1974) Oxygen control of reproductive growth: Possible mediation via dark respiration. *J. Exp. Bot.* **25**: 987-989.
- Gemel, J. and Randall, D.D. (1992) Light regulation of leaf mitochondrial pyruvate dehydrogenase complex: Role of photorespiratory carbon metabolism. *Plant Physiol.* **100**: 908-914.
- Goodman, H.M., Ecker, J.R. and Dean, C. (1995) The genome of *Arabidopsis thaliana*. *Proc. Nat'l. Acad. Sci. USA* **92**: 10831-10835.
- Grof, C.P.L., Winning, B.M., Scaysbrook, T.P., Hill, S.A. and Leaver, C.J. (1995) Mitochondrial pyruvate dehydrogenase: Molecular cloning of the E1 $\alpha$  subunit and expression analysis. *Plant Physiol.* **108**: 1623-1629.

- Harris, R.A., Popov, K.M., Shimomura, Y., Zhao, Y., Jaskiewicz, J., Nanaumi, N. and Suzuki, M. (1992) Purification, characterization, regulation and molecular cloning of mitochondrial protein kinase. *Adv. Enzyme Regul.* **32**: 267-284.
- Hill, R.L. and Bradshaw, R.A. (1969) Fumarase. *In: Methods in Enzymology*. Vol. XIII, J.M. Lowenstein (Ed.), Academic Press, New York. pp. 91-99.
- Hill, S.A., Grof, C.P.L., Bryce, J.H. and Leaver, C.J. (1992) Regulation of mitochondrial function and biogenesis in cucumber (*Cucumis sativus* L.) cotyledons during early seedling growth. *Plant Physiol.* **99**: 60-66.
- Hitz, W.D., Mauvis, C.J., Ripp, K.G., Reiter, R.J., DeBonte, L. and Chen, Z. (1995) The use of cloned rapeseed genes for cytoplasmic fatty acid desaturases and the plastid acyl-ACP thioesterases to alter relative levels of polyunsaturated and saturated fatty acids in rapeseed oil. *Proc. 9th Internat'nal Cambridge Rapeseed Congress UK*, pp. 470-472.
- Jefferson, R.A., Kavanaugh, T.A. and Bevan, M.W. (1987) GUS fusions:  $\beta$ -glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO J.* **6**: 3901-2907.
- Jorgensen, R.A. and Napoli, C.A. (1994) Genetic engineering of novel plant phenotypes. U.S. Patent No. 5283184.
- Katavic, V., Haughn, G.W., Reed, D., Martin, M. and Kunst, L. (1994) *In planta* transformation of *Arabidopsis thaliana*. *Mol. Gen. Genet.* **245**: 363-370.
- Katavic, V., Reed, D.W., Taylor, D.C., Giblin, E.M., Barton, D.L., Zou, J-T., MacKenzie, S.L., Covello, P.S. and Kunst, L. (1995) Alteration of fatty acid composition of an EMS-induced mutation in *Arabidopsis thaliana* affecting diacylglycerol acyltransferase activity. *Plant Physiol.* **108**: 399-409.

- Kinney, A.J. (1995) Improving soybean seed quality. *In: Induced Mutations and Molecular Techniques for Crop Improvement*. International Atomic Energy Agency, Vienna, Austria., pp. 101-113.
- Kinney, A.J. (1997) Genetic engineering of oilseeds for desired traits. *In: Genetic Engineering*, Vol. 19 (J.K. Setlow, ed.), Plenum press, NY., pp. 149-166.
- Kishore G.M. and Somerville, C.R. (1993) Genetic engineering of commercially useful biosynthetic pathways in transgenic plants. *Current Opinion in Biotechnology*. **4**: 152-158.
- Kleinig, H. and Liedvogel, B. (1980) Fatty acid synthesis by isolated chromoplasts from the daffodil. Energy sources and distribution patterns of acids. *Planta*, **150**: 166-169.
- Knutzon, D.S., Thompson, G.A., Radke, S.E., Johnson, W.B., Knauf, V.C., and Kridl, J.C. (1992) Modification of Brassica seed oil by anti-sense expression of a stearyl-acyl carrier protein desaturase gene. *Proc. Nat'l Acad. Sci. USA*, **89**: 2624-2628.
- Koncz, C. and Schell, J. (1986) The promoter of T<sub>L</sub>-DNA gene 5 controls the tissue-specific expression of chimaeric genes by a novel type of *Agrobacterium* binary vector. *Mol. Gen. Genet.* **204**: 383-396.
- Korotchkina, L.G. and Patel, M.S. (1995) Mutagenesis studies of the phosphorylation sites of recombinant human pyruvate dehydrogenase, *J. Biol. Chem.* **270**:14297-14304.
- Lagercrantz, U., Putterill, J., Coupland, G. and Lydiate, D. (1996) Comparative mapping in *Arabidopsis* and *Brassica*, fine scale genome collinearity and congruence of genes controlling flowering. *Plant J.* **9**: 13-20.
- Landschütze, V., Willmitzer, L. and Müller-Röber, B. (1995) Inhibition of flower formation by antisense repression of mitochondrial citrate synthase in transgenic potato plants leads to a specific disintegration of the ovary tissues of flowers. *EMBO J.* **14**: 660-666.

- Lassner, M.W., Lardizabal, K. and Metz, J.G. (1996) A jojoba  $\beta$ -ketoacyl-CoA synthase cDNA complements the canola fatty acid elongation mutation in transgenic plants. *The Plant Cell*, **8**: 281-292.
- Liedvogel, B. (1986) Acetyl Coenzyme A and isopentenyl diphosphate as lipid precursors in plant Cells-biosynthesis and compartmentation. *J. Plant Physiol.* **124**: 211-222.
- Liu, S., Baker, J.C. and Roche, T.E. (1995) Binding of the pyruvate dehydrogenase kinase to recombinant constructs containing the inner lipoyl domain of the dihydrolipoyl acetyltransferase component. *J. Biol. Chem.* **270**: 793-800.
- Lloyd, A.M., Walbot, V. and Davis, R.W. (1992) *Arabidopsis* and *Nicotiana* anthocyanin production activated by maize regulators *R* and *C1*. *Science* **258**: 1773-1775.
- Lutcke, H.A., Chow, K.C., Mickel, F.C., Moss, K.A., Kern, H.F. and Scheele, G.A. (1987) Selection of AUG initiation codon differs in plants and animals. *EMBO J.* **6**: 43-48.
- MacKenzie, S.L. and Jain, R.K. (1997) Improvement of oils crops via biotechnology. *Recent Res. Dev. In Oil Chem.* **1**: 149-158.
- Meyer, P. (1995) Understanding and controlling transgene expression. *Trends in Biotechnology*, **13**: 332-337.
- Meyerowitz, E.M. (1987) *Arabidopsis thaliana*. *Ann. Rev. Genet.* **21**: 93-111.
- Meyerowitz, E.M. and Chang, C. (1985) Molecular biology of plant growth and development: *Arabidopsis thaliana* as an experimental system. In: *Developmental Biology*, Vol. 5, Plenum Press, NY., pp. 353-366.
- Miernyk, J.A. and Randall, D.D. (1987) Some kinetic and regulatory properties of the pea mitochondrial pyruvate dehydrogenase complex. *Plant Physiol.* **83**: 306-310.
- Mol, J.M.N., Van der Krol, A.R., Van Tunen, A.J., Van Blokland, R., De Lange, P. and Stuitje, A.R. (1990) Regulation of plant gene expression by antisense RNA. *FEBS Lett.* **268**: 427-430.

- Moloney, M.M., Walker, J.M. and Sharma, K.K. (1989) High efficiency transformation of *Brassica napus* using *Agrobacterium* vectors. *Plant Cell Rep.* **8**: 238-242.
- Mullen, R.T., Lee, M.S., Flynn, C.R. and Trelease, R.N. (1997) Diverse amino acid residues function within the type 1 peroxisomal targeting signal: Implications for the role of accessory residues upstream of the type 1 peroxisomal targeting signal. *Plant Physiol.* **115**: 881-889.
- Murphy, L. and Scarth, R. (1994) Vernalization response in spring oilseed rape (*Brassica napus* L.) cultivars. *Can. J. Plant Sci.* **74**: 275-277.
- Nehra, N.S., Chibbar, R.N., Leung, N., Caswell, K., Mallard, C., Steinhauer, L. Baga, M. and Kartha K.K. (1994) Self-fertile transgenic wheat plants regenerated from isolated scutellar tissues following microprojectile bombardment with two distinct gene constructs. *Plant J.* **5**: 285-297.
- Nobukuni, Y., Mitsubuchi, H., Endo, F., Asaka, J., Oyama, R., Titani, K. and Matsuda, I. (1990) Isolation and characterization of a complementary DNA clone coding for the E<sub>1</sub> $\beta$  subunit of the bovine branched-chain  $\alpha$ -ketoacid dehydrogenase complex: Complete amino acid sequence of the precursor protein and its proteolytic processing. *Biochemistry* **29**: 1154-1160.
- Ohlrogge, J. and Browse, J. (1995) Lipid biosynthesis. *The Plant Cell*, **7**: 957-970.
- Okuley, J. Lightner, J., Feldmann, K., Yadav, N., Lark, E. and Browse, J. (1994) Arabidopsis *fad2* gene encodes the enzyme that is essential for polyunsaturated lipid synthesis. *The Plant Cell* **6**: 147-158.
- Padgett, S.R., Gruys, K.J., Mitsky, T.A., Tran, M., Taylor, N.B., Slater, S.C. and Kishore, G.M. (1997) Strategies for production of polyhydroxyalkanoate polymers in plants. *Plant Physiol. Suppl.*, **114**: 3 (abstract 10003).

- Parkinson, J.S. and Kofoed, E.C. (1992) Communication modules in bacterial signaling proteins. *Annu. Rev. Genet.* **26**: 71-112.
- Pearson, W.R. and Lipman, D.J. (1988) Improved tools for biological sequence comparison. *Proc. Natl. Acad. Sci. USA* **85**: 2444-2448.
- Popov, K.M., Zhao, Y., Shimomura, Y., Kuntz, M.J. and Harris, R.A. (1992) Branched-chain  $\alpha$ -ketoacid dehydrogenase kinase: Molecular cloning, expression and sequence similarity with histidine protein kinase. *J. Biol. Chem.* **267**: 13127-13130.
- Popov, K.M., Kedishvili, N.Y., Zhao, Y., Shimomura, Y., Crabb, D.W. and Harris, R.A. (1993) Primary structure of pyruvate dehydrogenase kinase establishes a new family of eukaryotic protein kinases. *J. Biol. Chem.* **268**: 26602-26606.
- Popov, K.M., Kedishvili, N.Y., Zhao, Y., Gudi, R. and Harris, R.A. (1994) Molecular cloning of the p45 subunit of pyruvate dehydrogenase kinase *J. Biol. Chem.* **269**: 29720-29724.
- Potrykus, I. (1991) Gene transfer to plants: Assessment of published approaches and results. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **42**: 205-225.
- Randall, D.D., Miernyk, J.A., Fang, T.K., Budde, R.J.A. and Schuller, K.A. (1989) Regulation of the pyruvate dehydrogenase complexes in plants. *Ann. NY Acad. Sci.* **573**: 192-205.
- Reid, E.E., Thompson, P., Lyttle, C.R. and Dennis, D.T. (1977) Pyruvate dehydrogenase complex from higher plant mitochondria and proplastids. *Plant Physiol.* **59**: 842-848.
- Rhodes, C.A., Pierce, D.A., Mettler, I.J., Mascarenhas, D. and Detmer, J.J. (1988) Genetically transformed maize plants from protoplasts. *Science* **240**: 204-207.
- Rosie, D. and Schatz, G. (1988) Mitochondrial presequences. *J. Biol. Chem.* **263**: 4509-4511.

- Roughan, P.G. and Ohlrogge, J.B. (1994) On the assay of acetyl-CoA synthetase in chloroplasts and leaf extracts. *Anal. Biochem.* **216**: 77-82.
- Roughan, P.G. and Ohlrogge, J.B. (1996) Evidence that isolated chloroplasts contain an integrated lipid-synthesizing assembly that channels acetate into long-chain fatty acids. *Plant Physiol.* **110**: 1239-1247.
- Sambrook, J. Fritsch, E.F. and Maniatis, T. (1989) Molecular Cloning, A Laboratory Manual, 2<sup>nd</sup>. Edition. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Sanford, J.C., Klein, T.M., Wolf, E.D. and Allen, N. (1987) Delivery of substances into cells and tissues using a particle bombardment process. *J. Part. Sci. Technol.* **5**: 27-37.
- Shimamoto, K., Terada, R., Izawa, T. and Fujimoto, H. (1989) Fertile transgenic rice plants regenerated from transformed protoplasts. *Nature* **338**: 274-276.
- Sinclair, T.R., Ward, J.P. and Randall, C.A. (1987) Soybean seed growth in response to long-term exposures to different oxygen partial pressures. *Plant Physiol.* **83**: 467-468.
- Somerville, C.R. (1993) Future prospects for genetic modification of the composition of edible oils from higher plants. *Am. J. Clin. Nutr.* **58** (2 Suppl.): 270S-275S.
- Songstad, D.D., Somers, D.A. and Griesbach, R.J. (1995) Advances in alternative DNA delivery techniques. *Plant Cell, Tissue and Organ Culture* **40**: 1-15.
- Southern E.M. (1975) Detection of specific sequences among DNA fragments separated by gele electrophoresis. *J. Mol. Biol.* **98**: 503-517.
- Srere, P.A. (1969) Citrate synthase. *In*: Methods in Enzymology. Vol. XIII, J.M. Lowenstein (Ed.), Academic Press, New York. pp. 3-11.

- Stark, D.M., Timmerman, K.P., Barry, G.F., Preiss, J. and Kishore, G.M. (1992) Regulation of the amount of starch in plant tissues by ADP glucose pyrophosphorylase. *Science* **258**: 287-292.
- Taylor, D.C., Magus, J.R., Bhella, R., Zou, J-T., MacKenzie, S.L., Giblin, E.M., Pass, E.W. and Crosby, W.L. (1992a) Biosynthesis of triacylglycerols in *Brassica napus* L. cv Reston; Target: Trierucin. *In: Seed Oils for the Future*. S.L. MacKenzie and D.C. Taylor (Eds.), American Oil Chemists' Society, Campaign, IL., pp. 77-102.
- Taylor, D.C., Barton, D.L., Rioux, K.P., Reed, D.W., Underhill, E.W., MacKenzie, S.L., Pomeroy, M.K. and weber, N. (1992b) Biosynthesis of acyl lipids containing very long-chain fatty acids in microspore-derived and zygotic embryos of *Brassica napus* L. cv Reston. *Plant Physiol.* **99**: 1609-1618.
- Taylor, D.C., Barton, D.L., Giblin, E.M., MacKenzie, S.L., Van den Berg, C.G.J. and McVetty, P.B.E. (1995) Developing seeds of a *Brassica oleracea* breeding line possess a lyso-phosphatidic acid acyltransferase capable of utilizing erucoyl-CoA and accumulating triacylglycerols containing erucic acid in the *sn*-2 position. *Plant Physiol.* **109**: 409-420.
- Vasil, I.K. (1994) Molecular improvement of cereals. *Plant Mol. Biol.* **25**: 925-937.
- Veeger, C., Der Vartanian, D.V. and Zeylemaker, W.P. (1969) Succinate dehydrogenase. *In: Methods in Enzymology*. Vol. XIII, J.M. Lowenstein (Ed.), Academic Press, New York. pp. 81-90.
- Voelker, T.A., Worrell, A.C., Anderson, L., Bleibaum, J., Fan, C., Hawkins, D.J., Radke, S.E., and Davies, H.M. (1992) Fatty acid biosynthesis redirected to medium chains in transgenic oilseed plants. *Science* **257**: 72-74.
- Voelker, T.A., Hayes, T.R., Cramner, A.M., Turner, J.C., and Davies, H.M. (1996) Genetic engineering of a quantitative trait: metabolic and



genetic parameters influencing the accumulation of laurate in rapeseed. *The Plant Journal* **9**: 229-241.

Walden, R. and Wingender, R. (1995) Gene-transfer and plant regeneration techniques. *Trends in Biotechnology* **13**: 324-331.

Zhang, H. Goodman, H.M. and Jansson, S. (1997) Antisense inhibition of photosystem I antenna protein Lhca4 in *Arabidopsis thaliana*. *Plant Physiol.* **115**: 1525-1531.

Zou, J-T. and Taylor, D.C. (1994) Isolation of an *Arabidopsis thaliana* cDNA homologous to parsley (*Petroselinum crispum*) S-adenosyl-L-methionine:trans-caffeoyl-Coenzyme A 3-O-methyltransferase, an enzyme involved in disease resistance. *Plant Physiol. Biochem.* **32**: 423-427.

Zou, J-T., Katavic, V., Giblin, E.M., Barton, D.L., MacKenzie, S.L., Keller, W.A., Hu, X. and Taylor, D.C. (1997) Modification of seed oil content and acyl composition in the Brassicaceae by expression of a yeast *sn-2* acyltransferase gene. *The Plant Cell* **9**: 909-923.

The teachings of the above references are hereby incorporated by reference.

CLAIMS:

1. Isolated and purified deoxyribonucleic acid (DNA), characterized in that said DNA includes a sequence according to SEQ ID NO:1 (ATCC 209562), or a part of SEQ ID NO:1, or a sequence that is substantially homologous to SEQ ID NO:1.
2. A vector for transformation of plant cells, characterized in that said vector contains a deoxyribonucleic acid sequence according to SEQ ID NO:1, or a part of SEQ ID NO:1, or a sequence that is substantially homologous to SEQ ID NO:1.
3. A vector according to claim 2, characterized in that said sequence is present in said vector in an anti-sense orientation.
4. A vector according to claim 2, characterized in that said sequence is present in said vector in a sense orientation.
5. Plasmid pYA5 (ATCC 209562).
6. Plasmid pAsYA5 (ATCC 209561).
7. A plant having a genome, characterized in that the genome contains an introduced nucleotide sequence of SEQ ID NO:1, or a part of SEQ ID NO:1, or a sequence that is substantially homologous to SEQ ID NO:1.
8. A plant seed having a genome, characterized in that said genome contains an introduced nucleotide sequence of SEQ ID NO:1, or a part of SEQ ID NO:1, or a sequence that is substantially homologous to SEQ ID NO:1.


9. A genetically transformed plant, characterized in that said genome has been transformed by a vector according to claim 2.
10. A genetically transformed plant seed, characterized in that said seed has been transformed by a vector according to claim 2.
11. A plant as claimed in 9, characterized by exhibiting an altered respiration rate compared to a genomically-unmodified plant of the same genotype.
12. A plant seed as claimed in 10, characterized by exhibiting an altered respiration rate compared to a genomically-unmodified plant of the same genotype.
13. A plant as claimed in 9, characterized by exhibiting an altered seed oil content compared to a genomically-unmodified plant of the same genotype.
14. A plant seed as claimed in 10, characterized by exhibiting an altered seed oil content compared to a genomically-unmodified plant of the same genotype.
15. A plant as claimed in 9, characterized by exhibiting an altered flowering time compared to a genomically-unmodified plant of the same genotype.
16. A plant seed as claimed in 10, characterized by said seed producing a plant which exhibits an altered flowering time compared to a genomically-unmodified plant of the same genotype.

17. A plant as claimed in 9, characterized by exhibiting an enhanced resistance to cold temperatures compared to a genomically-unmodified plant of the same genotype.
18. A plant seed as claimed in 10, characterized by said seed producing a plant which exhibits an enhanced resistance to cold temperatures compared to a genomically-unmodified plant of the same genotype.
19. A plant as claimed in 9, characterized by exhibiting an enhanced biomass compared to a genomically-unmodified plant of the same genotype.
20. A plant seed as claimed in 10, characterized by said seed producing a plant which exhibits an enhanced biomass compared to a genomically-unmodified plant of the same genotype.
21. A plant as claimed in 9, characterized by exhibiting an enhanced capacity to accumulate biopolymers compared to a genomically-unmodified plant of the same genotype.
22. A plant seed as claimed in 10, characterized by said seed producing a plant which exhibits an enhanced capacity to accumulate biopolymers compared to a genomically-unmodified plant of the same genotype.
23. A method of producing transgenic plants by introducing a nucleotide sequence into a genome of said plant, characterized in that said nucleotide sequence introduced into said genome includes SEQ ID NO:1, or a part of SEQ ID NO:1, or a sequence that is substantially homologous to SEQ ID NO:1.

24. A method according to claim 23, characterized in that said plant is a member of the *Brassicaceae*.
25. A method according to claim 23, characterized in that said plant is a member of the group consisting of borage (*Borago* spp.), Canola, castor (*Ricinus communis*), cocoa bean (*Theobroma cacao*), corn (*Zea mays*), cotton (*Gossypium* spp), *Crambe* spp., *Cuphea* spp., flax (*Linum* spp.), *Lesquerella* and *Limnanthus* spp., Linola, nasturtium (*Tropaeolum* spp.), *Oenothera* spp., olive (*Olea* spp.), palm (*Elaeis* spp.), peanut (*Arachis* spp.), rapeseed, safflower (*Carthamus* spp.), soybean (*Glycine* and *Soja* spp.), sunflower (*Helianthus* spp.), tobacco (*Nicotiana* spp.), *Vernonia* spp., wheat (*Triticum* spp.), barley (*Hordeum* spp.), rice (*Oryza* spp.), oat (*Avena* spp.) sorghum (*Sorghum* spp.), rye (*Secale* spp.) or other members of the *Gramineae*.
26. A plant DNA sequence or part thereof, characterized in that the sequence is substantially homologous to at least a part of SEQ ID NO:1, and in that said sequence has been isolated or characterized using sequence information from SEQ ID NO:1.
27. A method of changing the oil content of plant seeds by introducing a sense or anti-sense nucleic acid construct into a plant transformation vector, using the vector to transform the genome of a plant or plant seed, and then growing the plant or plant seed and extracting the oil from the plant seed, characterized in that said nucleic acid sequence is SEQ ID NO:1, or a part of SEQ ID NO:1, or a sequence that is substantially homologous to SEQ ID NO:1.
28. A method of changing the biopolymer content of a plant or plant storage organ by introducing an anti-sense or sense construct into a plant

transformation vector, using the vector to transform the genome of a plant or plant storage organ, and then growing the plant or plant storage organ and extracting the biopolymer from the plant or plant storage organ, characterized in that the construct contains SEQ ID NO:1, or a part of SEQ ID NO:1, or a sequence that is substantially homologous to SEQ ID NO:1.

1/14

**FIG. 1**

2/14

## Figure 2

The nucleotide sequence and its deduced amino acid sequence of the *Arabidopsis thaliana* pyruvate dehydrogenase kinase (PDH kinase; PDHK) cDNA (clone YA5; ATCC # 209562).

```

1: t cca tct gcg cac ttc ttt cgt cca gtc gat gat aat aac ggt gg
46: a gaa cga cgg agg cgg gcg acg tta ggg ttt cta atc att tct ct
91: c tct tag acg ctt atg gca gtg aag aaa gcc tgc gaa atg ttc cc
      M  A  V  K  K  A  C  E  M  F  P

136: g aag agt ttg atc gaa gat gtt cac aaa tgg ggt tgc atg aag ca
      K  S  L  I  E  D  V  H  K  W  G  C  M  K  Q

181: a acc ggt gtt agc ctt aga tac atg atg gag ttt ggt tcc aaa cc
      T  G  V  S  L  R  Y  M  M  E  F  G  S  K  P

226: t act gag agg aat ctt ttg att tct gct cag ttt ttg cat aag ga
      T  E  R  N  L  L  I  S  A  Q  F  L  H  K  E

271: g ctt ccg att cgc gtc gcc agg aga gcg atc gaa ctc cag acg ct
      L  P  I  R  V  A  R  R  A  I  E  L  Q  T  L

316: t cct tat ggt ctc tct gat aaa cct gcc gtt ttg aag gtg cgg ga
      P  Y  G  L  S  D  K  P  A  V  L  K  V  R  D

361: t tgg tat ttg gaa tct ttc agg gac atg aga gca ttt cct gag at
      W  Y  L  E  S  F  R  D  M  R  A  F  P  E  I

406: t aag gat tcg ggt gac gag aaa gat ttc act cag atg att aag gc
      K  D  S  G  D  E  K  D  F  T  Q  M  I  K  A

451: t gtc aaa gta agg cat aac aat gtg gtt ccc atg atg gct ttg gg
      V  K  V  R  H  N  N  V  V  P  M  M  A  L  G

496: t gtt aat cag ctc aag aaa gga atg aat tct gga aat ctt gat ga
      V  N  Q  L  K  K  G  M  N  S  G  N  L  D  E

541: g att cat cag ttt ctt gat cgt ttc tac ttg tcg cga atc ggg at
      I  H  Q  F  L  D  R  F  Y  L  S  R  I  G  I

586: c cgg atg ctt att ggg cag cac gtt gag ttg cat aat cca aat cc
      R  M  L  I  G  Q  H  V  E  L  H  N  P  N  P

631: a ccg ctt cat aca gtg ggt tat ata cac aca aag atg tct cct at
      P  L  H  T  V  G  Y  I  H  T  K  M  S  P  M

676: g gag gta gca agg aat gca agt gaa gat gct cgg tca att tgt tt
      E  V  A  R  N  A  S  E  D  A  R  S  I  C  F

721: c cga gag tac ggt tct gca ccg gaa ata aac ata tat ggc gat cc
      R  E  Y  G  S  A  P  E  I  N  I  Y  G  D  P

766: c agt ttc acc ttt ccg tat gtt cca acg cat ttg gat ctt atg at
      S  F  T  F  P  Y  V  P  T  H  L  D  L  M  M

811: g tat gag cta gtc aag aac tct cta cgt gct gtc caa gag cga tt
      Y  E  L  V  K  N  S  L  R  A  V  Q  E  R  F

856: t gtt gac tct gat aga gtt gca cca cca atc cgc att ata gtt gc
      V  D  S  D  R  V  A  P  P  I  R  I  I  V  A

```



3/14

901: t gat gga atc gaa gat gtt act ata aag gtc tca gat gaa ggt gg  
D G I E D V T I K V S D E G G

946: a ggt ata gca aga agc ggt ctt ccc aga ata ttc acc tat ctt ta  
G I A R S G L P R I F T Y L Y

991: c agc act gca aga aac ccg ctt gag gag gat gtc gat tta gga at  
S T A R N P L E E D V D L G I

1036: a gct gat gtt ccc ggg act atg ggt gga tat ggt tat ggt ctt cc  
A D V P G T M G G Y G Y G L P

1081: a att agt cgc ttg tat gct cga tat ttc ggt gga gat ttg cag at  
I S R L Y A R Y F G G D L Q I

1126: c ata tcc atg gaa gga tat ggg act gat gca tac ttg cac ttg tc  
I S M E G Y G T D A Y L H L S

1171: t cgc ctt gga gat tcg caa gag cct tta ccc tga gaa cat ctc ta  
R L G D S Q E P L P \*

1216: t gtc agg caa agt aaa gaa agc ttt gac atg tat tta tgg tag at

1261: g agg gat atc tac aat act caa tta ttt atg ctt ttc cag ttt ct

1306: g cta atg tac aga cta cag aca tta ttt tct cgt att acg ctt tc

1351: t tga ttt tag act cag ata tgg agc ttt ttc caa gtg agt taa tc

1396: t cct atg att tgt ttt ggt tcg atc caa aac cac ctt gta tcc ga

1441: a aaa aaa aaa aaa aaa a

4/14

Figure 3

The amino acid sequence alignment of the *Arabidopsis* PDH kinase (Ya5p) with other mammalian mitochondrial ketoacid dehydrogenase kinase. **Pdhk I**, porcine PDH kinase subunit I. **Pdhk II**, porcine PDH kinase subunit II. **Bckdhk**, porcine branched chain alpha-ketoacid dehydrogenase kinase. Dots indicate gaps. Identical amino acid residues are highlighted in bold upper case type.

Ya5p	MAVKKA.....CEMFPK
pdhkI	MrwFRAlIknaSlagapkyiehfskfsp.....spLsmK
pdhkII	MrLaRllrggtsvrplcavpcasrslasdsasgsgpasesgvpgqvdfyarfspLsmK
bckdhk	MiLtsvlsgsprsgsslwpllgsslsrlvrstsatdthhvelare.....rsktvt
ya5p	SLIEDVHKWGCmkQTGVSLR..YMMEFGSKPTERNLLISAQFLHKELPIRVARRAIELQT
pdhkI	qfLDfgssnACeKtSftfLRqelpVrLAnimkEiNLLpdrvlstpsVqLvqSwyvqsLld
pdhkII	qfLDfgsvnACeKtsfMfLRqelpVrLAnimkEisLLpdnllrtpsVqLvqSwyiqsLQe
bckdhk	SfyNgsaidvvaekPSVrLtptmMLysGrsqdgshLLkSGrYLqQELPVRIhRikgfvv
ya5p	LPYGLSDKPAVLKVRDWYLESFRDMRAFPEIKDSGDEKDFtQMIKAVKVRHNNVPMAL
pdhkI	ImefLdkdPedhRtlSqFtDALvtIRn.....RHNNVVPtMAq
pdhkII	LldfkdkSAedaKtiyeFtDTvirIRn.....RHNDVIPtMAq
bckdhk	flsSLvatlPyctVhELyIRAFqkLtdFPpIKDqADEaQYcQLVRqLlddHkDVvtLLAe
ya5p	GVNQLKKGmNSGNLDEIH.Q.FLDRFYLSRIGIRMLIGQHVLEHNPPLHTVGyI...H
pdhkI	GVLEyKdTygddpVsNqniQyFLDRFYLSRISIRMLInQtLifdgStnPaHPkhgsidpn
pdhkII	GVNEyKeSfgSdpVtsqnvQyFLDRFYMSRISIRMLLnQHsLlfgGkGSpsrhkhIgsin
bckdhk	GlrEsRKhIEdekL..Vry..FLDKtltSRLGIRMLaThHLaLH..EdkpdfVGiIs...
ya5p	TKMSPMEVARNASEDARSICFREYGSAPeINIYGDPS.....FTFPYVPTHLdLMMYEL
pdhkI	PncSvsDVvKDayDmAkLLCdKyYmASPDLEIqevnAtnatqpihmVYVPSHlyhMLFEL
pdhkII	PncdvvEViKDGyENARrLCdlyYvnSPELEleelnAkspgqpiqvVYVPSHlyhMVfEL
bckdhk	TRLSPkkIiekwvDfARrLCeHkYGNAPrVrInGhvAa.....rFPFIPmpLDyILpEL
ya5p	VKNSLRAVQERFVDSDRVAPPRIIIVADGIEDVTIKVSDEGGGIARSGLP...RIFTYL
pdhkI	fKNAMRatvEshesSlt.lPPIkIMVALGeEDLSIKMSDrGGGVPlrkIe...RLFSYM
pdhkII	fKNAMRatmEhhaDKgv.yPPIqvhVtlGeEDLTVKMSDrGGGVPlrkId...RLFnYM
bckdhk	LKNAMRatmEshLDtpynvPdVvItIANndvDLiIrISDrGGGIAHkdLdrvmDyHFTta
ya5p	YSTARNPLEEDVDLGIADVPGTMG.....GYGYGLPISRLYARYFGGDLQIISMEGY
pdhkI	YSTApTqpgtg.gTp.....LA.....GFGYGLPISRLYAKYFQGDQLQFSMEGf
pdhkII	YSTAprPrvEtsravp.....LA.....GFGYGLPISRLYAqYFqGDkLySLEGY
bckdhk	....eaStQDprisPLfDhldThSggqsgpmhGFGFGLPtSRaYAeYlGGsLQLqSLQGi
ya5p	GTDAYLHLSRLGDSQEPLP
pdhkI	GTDAviyLkaLSTdSverlpvynksawrhYqtigeagdwcvpstepkntyrvs
pdhkII	GTDAviyikaLSTeSierlpvynkaawkhYrtnheaddwcvpsrepkdmTtfrrs
bckdhk	GTdv.LHrSRhidgreesfri

5/14






FIG. 4

6/14

**FIG. 5**

7/14



Northern Blot Analysis of PDK Gene Expression in *Arabidopsis thaliana*  
RNA was extracted from flowers (F), seedlings(L), young siliques(YS)  
and maturing siliques (MS).

**FIG. 6**

8/14



YA5 and control  
E.coli lysate, SDS PAGE

**FIG. 7a**




Human E1 $\alpha/\beta$ , SDS PAGE  
Obtained from Dr. M.S. Patel

**FIG. 7b**

Phosphorylation of PDHE1 subunit:

YA5 *E.coli* lysate, Human E1 $\alpha/\beta$ ,  $\gamma$ -<sup>32</sup>P-ATP, Mg<sup>++</sup>, pH 7.0




**FIG. 7c**

**FIG. 7d**


9/14

Leaf Mitochondrial PDC Activity in *A. thaliana* Non-Transformed WT and T<sub>2</sub> Transgenic Plants Containing Constitutively Expressed Anti-Sense PDHK

**FIG. 8**

10/14

Leaf Mitochondrial Citrate Synthase Activity in *A. thaliana* Non-Transformed WT and T<sub>2</sub> Transgenic Plants Containing Constitutively Expressed Anti-Sense PDHK




**FIG. 9**



11/14

Seed oil content ( $\mu\text{g}$  total fatty acids / 100 seeds) in *A. thaliana* non-transformed wild-type controls (n-WT Con) and the T<sub>2</sub> generation of pBI121 plasmid only control (pBI121 Con) and antisense PDHK (YA5) transgenics

**FIG.10**

12/14






FIG. 11

13/14

**FIG. 12**

14/14

Average Number of Rosette Leaves in *A.thaliana* non-transformed WT and T<sub>2</sub> pBI121 control and anti-sense PDHK transgenic plants upon entering the generative phase (n=5-6)

**FIG. 13**

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 98/00096

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/29 C12N15/63 A01H5/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	E.E. REID ET AL.: "Pyruvate dehydrogenase complex from higher plant mitochondria and proplastids" PLANT PHYSIOLOGY, vol. 59, 1977, pages 842-848, XP002070250 cited in the application *see the whole article* ---	1-28
A	C.P.L. GROF ET AL.: "Mitochondrial pyruvate dehydrogenase" PLANT PHYSIOLOGY, vol. 108, 1995, pages 1623-1629, XP002070251 cited in the application *see the whole article* ---	1-28
-/--		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "Z" document member of the same patent family

Date of the actual completion of the international search

3 July 1998

Date of mailing of the international search report

23/07/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Marie, A

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 98/00096

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>T. NEWMAN ET AL.: "Genes galore: a summary of methods for accessing results from large scale partial sequencing of anonymous Arabidopsis cDNA clones"</p> <p>PLANT PHYSIOLOGY, vol. 106, 1994, pages 1241-1255, XP002070252</p> <p>*see the whole article*</p> <p>-----</p>	1-28